PHILOSOPHICAL TRANSACTIONS,

OF THE

ROYAL SOCIETY

OF

LONDON.

FOR THE YEAR MDCCCXIII.

PART II.

LONDON,

PRINTED BY W. BULMER AND CO. CLEVELAND-ROW, ST. JAMES'S;
AND SOLD BY G. AND W. NICOL, PALL-MALL, BOOKSELLERS TO HIS MAJESTY,
AND PRINTERS TO THE ROYAL SOCIETY.

MDCCCXIII.

PHILOSOPHICAL TRANSACTIONS

CONTENTS.

XIX. *An Account of some organic Remains found near Brentford, Middlesex. By the late Mr.* William Kirby Trimmer. *Communicated in a Letter from Mr.* James R. Trimmer *to the Right Hon. Sir* Joseph Banks, *Bart. K. B. P. R. S.* p. 131

XX. *On a new Construction of a Condenser and Air-pump. By the Rev.* Gilbert Austin. *In a Letter to Sir* Humphry Davy, *LL.D. F. R. S.* p. 138

XXI. *On the Formation of Fat in the Intestines of living Animals. By Sir* Everard Home, *Bart. Presented by the Society for promoting the Knowledge of Animal Chemistry.* p. 146

XXII. *On the colouring Matter of the black Bronchial Glands, and of the black Spots of the Lungs. By* George Pearson, *M. D. F. R. S.* p. 159

XXIII. *Experiments on the Alcohol of Sulphur, or Sulphuret of Carbon. By* J. Berzelius, *M. D. F. R. S. Professor of Chemistry at Stockholm; and* Alexander Marcet, *M. D. F. R. S. one of the Physicians to Guy's Hospital.* p. 171

XXIV. *On the Means of procuring a steady Light in Coal Mines without the danger of Explosion. By* William Reid Clanny, *M. D. of Sunderland. Communicated by* William Allen, *Esq. F. R. S.* p. 200

XXV. *On the Light of the Cassegrainian Telescope, compared with that of the Gregorian. By Captain* Henry Kater, *Brigade-Major. Communicated by the Right Hon. Sir* Joseph Banks, *Bart. K. B. P. R. S.* p. 206

XXVI. *Additional Observations on the Effects of Magnesia in preventing an increased Formation of Uric Acid; with Remarks on the Influence of Acids upon the Composition of the Urine.* By William Thomas Brande, *Esq. F. R. S. Prof. Chem. R. I. Communicated by the Society for improving Animal Chemistry.* p. 213

XXVII. *Additions to an Account of the Anatomy of the Squalus Maximus, contained in a former Paper; with Observations on the Structure of the Branchial Artery.* By Sir Everard Home, *Bart. F. R. S.* p. 227

XXVIII. *Some further Observations on a new detonating Substance. In a Letter from* Sir Humphry Davy, *LL.D. F.R.S, V.P.R.I. to the Right Hon. Sir* Joseph Banks, *Bart. K.B. P.R.S.* p. 242

XXIX. *Experiments on the Production of Cold by the Evaporation of the Sulphuret of Carbon.* By Alexander Marcet, *M.D. F.R.S. one of the Physicians to Guy's Hospital.* p. 252

XXX. *On a saline Substance from Mount Vesuvius.* By James Smithson, *Esq. F. R. S.* p. 256

XXXI. *Some Experiments and Observations on the Substances produced in different chemical Processes on Fluor Spar.* By Sir Humphry Davy, *LL.D. F.R.S. V.P.R.I.* p. 263

XXXII. *Catalogue of North Polar Distances of Eighty-four principal fixed Stars, deduced from Observations made with the Mural Circle at the Royal Observatory.* By John Pond, *Esq. Astronomer Royal, F. R. S.* p. 280

XXXIII. *Observations of the Summer Solstice, 1813, with the Mural Circle at the Royal Observatory.* By John Pond, *Esq. Astronomer Royal, F. R. S.* p. 304

Presents received by the Royal Society, from November 1812 to July 1813. p. 305

Index.

XIX. *An Account of some organic Remains found near Brentford, Middlesex. By the late Mr.* William Kirby Trimmer. *Communicated in a Letter from Mr.* James R. Trimmer *to the Right Hon. Sir* Joseph Banks, *Bart. K. B. P. R. S.*

Read March 4, 1813.

Sir,

My late brother, in digging clay near Brentford, for the manufacture of bricks and tiles, frequently found the remains of animals and extraneous fossils, which you, Sir, with other Gentlemen of the Royal Society, very often inspected, and the interest which you took in them added much to his ardour in collecting them.

A few days before he was attacked with the illness which terminated in his death, he had drawn up an account of the different strata in which the organic remains in his cabinet were found deposited, with the intention of submitting it to you. His long illness and other circumstances have delayed my communicating to you earlier the enclosed letter, which I found amongst his papers; and I request, Sir, that if you

should deem it of sufficient interest, you will do me the honour of laying it before the Royal Society.

I have the honour to be, Sir,
with great respect,
your much obliged humble servant,

JAMES R. TRIMMER.

Brentford, Feb. 1813.

To the Right Hon. Sir Joseph Banks, *Bart. &c. &c. &c.*

SIR,

My collection of organic remains having been nearly formed under your own eye, and my zeal in the collection, and care in the preservation of them, having been greatly increased by the importance which yourself and other Gentlemen of the Royal Society have attached to them, I do myself the honour of presenting to you the following statement of the manner in which they were deposited in the strata where they have been found; conceiving that however curious the specimens may be in themselves, they can conduce but little to the advancement of geological knowledge, if that part of their history is wanting.

The specimens have been collected from two fields, not contiguous to each other; therefore to avoid confusion, I shall take each field separately, first describing the strata as far as they have come within my knowledge, and afterwards I shall speak of the organic remains as they were respectively found in those strata.

The first field is about half a mile north of the Thames at Kew bridge; its surface is about twenty-five feet above the

Thames at low water. The strata here are first, sandy loam from six to seven feet, the lowest two feet slightly calcareous. Second, sandy gravel a few inches only in thickness. Third, loam slightly calcareous from one to five feet; between this and the next stratum, peat frequently intervenes in small patches of only a few yards wide, and a few inches thick. Fourth, gravel containing water; this stratum varies from two to ten feet in thickness, and is always the deepest in the places covered by peat; in these places the lower part of the stratum becomes an heterogeneous mass of clay, sand, and gravel, and frequently exhales a disagreeable muddy smell. Fifth, the main stratum of blue clay, which lies under this, extends under London and its vicinity, the average depth of this clay has been ascertained, by wells that have been dug through it, to be about two hundred feet under the surface of the more level lands, and proportionably deeper under the hills, as appears from Lord SPENCER's well at Wimbledon, which is five hundred and sixty-seven feet deep. This stratum, besides figured fossils, contains pyrites and many detached nodules; at the depth of twenty feet there is a regular stratum of these nodules, some of which are of very considerable size.

In the first stratum, as far as my observation has extended, no remains of an organised body has ever been found, and as my search has not been very limited, I may venture to say it contains none. In the second stratum, snail shells, and the shells of river fish have been found, and a few bones of land animals, but of inconsiderable size, and in such a mutilated state, that it cannot be ascertained to what class they belong. In the third stratum, the horns and bones of the ox, and the horns, bones, and teeth of the deer have been found, and also, as

in the second stratum, snail shells and the shells of river fish. In the fourth stratum were found teeth and bones of both the African and Asiatic elephant, teeth of the hippopotamus, bones, horns, and teeth of the ox.

A tusk of an elephant measured, as it lay on the ground, nine feet three inches, but in attempting to remove it, it broke into small pieces. When this stratum dips into the clay, and becomes a mixed mass, as before stated, it is seldom without the remains of animals. In the fifth stratum, namely the blue clay, the extraneous fossils are entirely marine, with the exception of some specimens of fruit and pieces of petrified wood, the latter of which may be considered as marine, because when of sufficient size, they are always penetrated by teredines. The other fossils from this stratum are nautili, oysters, pinnæ marinæ, crabs, teeth and bones of fish, and a great variety of small marine shells; this stratum has been penetrated hitherto in this field only to the depth of thirty feet, throughout which the specimens found were dispersed, without any regularity.

The second field is about one mile to the westward of the former, one mile north of the Thames, and a quarter of a mile to the eastward of the river Brent; its height above the Thames, at low water, is about forty feet. The strata are, first, sandy loam, eight or nine feet, in the lowest three feet of which it is slightly calcareous. Second, sand, becoming coarser towards the lowest part, and ending in sandy gravel from three to eight feet. Third, sandy loam highly calcareous, having its upper surface nearly level, but gradually increasing in thickness from a feather edge to nine feet. Below this are two strata of gravel and clay, as in the other field, but as these strata have been only occasionally penc-

trated in digging for water, nothing therefore is known with respect to them, but that they exist there.

In the first stratum, as in the other field, no organic remains have been observed. In the second, but always within two feet of the third stratum, have been found the teeth and bones of the hippopotamus, the teeth and bones of the elephant, the horns, bones, and teeth of several species of deer, the horns, bones, and teeth of the ox, and the shells of river fish.

The remains of hippopotami are so extremely abundant, that in turning over an area of one hundred and twenty yards in the present season, parts of six tusks have been found of this animal, besides a tooth and part of the horn of a deer, part of a tusk, and part of a grinder of an elephant, and the horns with a small part of the skull of an ox. One of these horns I had an opportunity of measuring, as it lay on the ground, and found it to be four feet and a half in length, following the curve, and five inches in diameter at the large end; it was found impracticable to remove it, otherwise than in fragments, which I have preserved, and have hopes of being able to put a considerable part of it together. The immense size of this horn is rendered more remarkable, by another horn from the same spot, which measures but six inches in length. Though this stratum is so extremely productive of the remains of animals, yet there are but few good cabinet specimens from it, owing, it is presumed, to their having been crushed at the time they were buried, and to the injury they have since received from moisture. It is necessary to remark, that the gravel stones in this stratum do not appear to have been rounded in the usual way by attrition, and that the bones must have been deposited after the flesh was off,

because, in no instance have two bones been found together which were joined in the living animal; and further, that the bones are not in the least worn, as must have been the case had they been exposed to the wash of a sea beach.

In the third stratum, viz. calcareous loam, have been found the horns, bones, and teeth of the deer, the bones and teeth of the ox, together with snail shells and the shells of river fish.

Brentford, in the neighbourhood of which are the fields I have mentioned, is situated on the north bank of the Thames, and is six miles west of London.

The fall of the Thames from Brentford to its mouth at the Nore,* is estimated at seven feet.

I have the honour to be,
with great respect, Sir,
your much obliged
and obedient humble servant,

WM. KIRBY TRIMMER.

Brentford.

* Col. Mudge, Trigonometrical Survey, p. 85.

Fig. 1.



Fig. 2.









EXPLANATION OF THE PLATES.

The following sketches (see Plates VIII, IX, X, XI, XII.) will give a tolerably correct idea of the state in which the bones were found.

PLATE VIII.

Fig. 1. The grinding tooth of an Asiatic elephant.
Fig. 2. The grinding tooth of an African elephant.

PLATE IX.

Three grinding teeth of the hippopotamus.

PLATE X.

A tusk and fore tooth of the hippopotamus.

PLATE XI.

Bones of the deer kind. A lower jaw, two grinding teeth, and a bone of the leg.

PLATE XII.

Two portions of the horns of the deer. One set upon a bony pedicle, which is not the case in any recent deer, except in a small species found in Prince's Island in the East Indies.

XX. *On a new Construction of a Condenser and Air-pump. By the Rev.* Gilbert Austin. *In a Letter to Sir* Humphry Davy, *LL.D. F.R.S.*

Read March 11, 1813.

DEAR SIR,

Woodville, Dec. 5, 1811.

ACCORDING to your desire, I give you a description of my glass condenser, which, upon inspection, you were so obliging as to consider might possibly be found useful in some of those extensive and profound chemical researches, in which you are happily engaged.

In the year 1799, I sent to the Royal Irish Academy the description of a little condensing instrument for the extemporaneous preparation of artificial mineral waters, which I stated would be more proper for nice experiments, if it could be made entirely of glass. As this suggestion had not, to my knowledge, induced any one better qualified to construct such an instrument of that material; pursuing my own ideas, I contrived the glass condenser herein described, the different parts of which, as to their fittings and application, that is the grinding of the barrel, the fitting and connecting of the flat joints and valves, I have been obliged to execute myself.

I have found this instrument to answer my immediate purpose very well, and have also applied it successfully to some other trivial chemical experiments; but it has for a considerable time (on account of my other avocations) remained neglected, and would have continued so to remain, had not you

given me hopes that it might be admitted to a place in your laboratory.

It was my wish, in the construction of this instrument, to have made the piston of glass only, without any packing of flax or thread for fitting it to the barrel; but I have not ventured to attempt so much. And perhaps as the packing is only a very small ring of a known substance, and as that may be varied at pleasure, by using for the purpose silk, or wool, or leather, or elastic gum, the effect in any experiment may be easily appreciated. The water which lubricates the barrel, and with which the joints are wetted in the first instance only (for the permanent junctures require no second wetting, but remain perfect for any length of time) may require also to be taken into the account, and if on occasion it should be thought preferable, oil may be used in the place of water. Should it however appear to you, that the want of a piston of glass, which should act without the addition of any foreign substance, is an important defect, you may possibly get one executed perfectly by some of those ingenious and excellent artists who are to be found in London. You will probably also improve on the form and construction of this first instrument (as far as I know) made of glass for the purposes mentioned. I have myself partly executed another of more general application on similar principles, of which, with your kind permission, I shall give you an account, as soon as I have completed it.

The great impediment to the forming of pneumatic instruments of glass, appears to me to have arisen from the difficulty of making the necessary joinings, in such a manner as to be capable of sustaining very considerable pressure, to be easily

disunited or connected, as occasion might require, and to be perfectly air tight. The conical junctures hitherto used, do not bear much pressure, and are objectionable in other respects: I have therefore formed my joints of plane surfaces, and find them as perfect, as permanent, and as easy in application as I can desire. For the construction of these joints, it is necessary that the glass vessels, or pieces to be connected, should be so formed that the part where they are to be joined should admit of having a broad projecting ring cast upon it. The ring should be formed by the glass-blower, as truly square and flat as convenient. The face is then to be ground (but not polished) perfectly plane, and then it will of course fit any other plane surface;—a great convenience on many occasions, and an advantage peculiar to the flat joint.

The connecting pieces, or collars for the joints, are formed of a flat ring of brass for each, about half an inch broad, and above one-eighth of an inch thick; this ring must be just wide enough to pass over the glass ring to which it is adapted. Fig. 3. *a* (Pl. XIV.) A mahogany ring about half an inch thick, having the diameter of the hole equal to that of the neck of the vessel above the glass ring, is to be screwed to the brass, and then sawed into two pieces through its diameter, fig. 3, *b*. These pieces are to be unscrewed, and applied to the neck, and they are then to be screwed to their places on the brass. The compound ring, or collar, will be thus secured above the glass ring, and may be fastened to another collar, attached in the same manner to the piece to be joined to it, by four screws passing through both when the joint is intended to be in some degree permanent, as in fig. 3, *c*. in which (*cc*) represents the glass pieces to be joined, (*aa*) the brass ring, and (*bb*) the

ring of mahogany. But if the joint require to be frequently separated in using the instrument, it may be applied moistened to the other, and pressed closely and perfectly by three milled-headed screws in the brass bridge, which bear upon the collar or ring of brass with their points. The degree of pressure necessary is not to be very great, but it must be equably exerted and felt on each screw, in order that the surfaces should be applied truly, and that the joint may be air tight—a very little practice will render this operation easy and certain. The bridge with its screws, &c. is represented, fig. 4, *a* (Pl. XIV.)—The plan of the bridge, and the end of the barrel with the valve in the piston, is seen at (*b*).—(*c*) is a stop to prevent the piece to be joined to the barrel from being pushed too far.

The glass valves are formed of plane convex lenses not polished, fitted to hollow spherical cavities, as in figs. 5 and 6.

These particulars being premised, the description of the instrument will be more easily understood.

Fig. 1, (Pl. XIII.) Represents a perspective view of the glass condenser, mounted on its frame and stand.

Fig. 2, (Pl. XIII.) A side view of the glass pieces of the condenser, disencumbered of the frame and connecting collars. The piston is seen outside of the barrel, and the iron sliding bar, with its arms which support the barrel, is represented with its clamps—but without the mahogany pieces. The same letters refer to the same parts of the instrument in figs. 1 and 2.

In figure 2 are also represented the valves in their places—the joints connected by dotted lines are permanent.

Figs. 1 and 2.

(*a*) The upper vessel which contains the liquid to be impregnated with the gas. A safety valve closes its mouth, seen at large, fig. 5 (Pl. XIV.)

(*b*) A piece connected permanently to the upper vessel; it carries the valve, which is so far below its surface as to allow it to rise a little in action, but not to turn over: the plane side of the lens is indented with a deep cross cut, to allow the gas to pass over it into the upper vessel, seen at large, fig. 6, *g*. *d* (Pl. XIV.)

(*c*) The glass barrel having a broad ring on each end.

Length of the barrel	-	13. inches.
Diameter of the bore	- -	0.$\frac{7}{10}$
External diameter	-	1.$\frac{3}{4}$
Diameter of the broad rings	-	2.$\frac{1}{4}$

(*d*) The hollow glass piston having one broad ring next (*e*), for the flat joint which connects the piston with it, and another ring, about four inches above this last, in order to secure it the better to the frame in which this ring and the piston are more than half sunk, and then covered over by a piece of mahogany pressing upon it with four screws. The covered part of the piston is represented by the dotted lines near (*d*) in fig. 1. At the end of the piston are two small rings, to confine the packing by which it is fitted to the barrel; and the extremity is ground into the hollow segment of a sphere, into which is fitted an unpolished plane convex lens, fig. 2, and seen at large, fig. 6.

(*e*) The receiver, from which the gas is drawn up and condensed into the upper vessel (*a*). It is furnished with a ground stopper, which may be used occasionally as hereafter mentioned.

(*f*) The pneumatic trough placed on a stool under the receiver.

(*g*) A retort, in which the gas is generated as usual.

(*hh*) Two pieces of mahogany, touching the sides of the barrel, and in their length fitting between its rings, so as when raised or depressed, to move the barrel without shake, and supporting the platform (*i*) with its bridge, as seen at large, fig. 4. These pieces are held together by iron clamps hollowed, so as to admit the barrel freely, fig. 2, *k*: they are clamped behind in the same manner, and then the barrel is secured by wedges in its proper situation, so as to be fixed perfectly parallel to the iron sliding bar. From the clamps behind proceed two strong arms (*kk*), fixed at right angles in the sliding bar (*l*), which moves equably in the long brass dove-tailed groove (*m*). This dove-tailed groove is screwed to the strong upright pillar (*o*). By means of the handle (*n*), which is fastened to the sliding bar between the two strong arms (*kk*), the barrel, with its apparatus, the platform (*i*), the brass bridge confining the upper vessel (*a*), by the piece (*b*), are alternately raised and depressed. In this action, the barrel moves upon the fixed hollow piston (*d*). The gas is extracted from the receiver at every ascending stroke, and passing through the hollow glass piston (*d*) above its valve, which opens upwards, is forced at every descending stroke through the valve between (*a* and *b*) into the upper vessel containing the liquor to be impregnated. Should more gas be forced into the vessel than the liquid readily absorbs, the safety valve at the top (seen at large, fig. 5) allows it to escape.

Fig. 5, (Pl. XIV.) is an enlarged view of the mouth of the upper vessel (*a*), with the collar (*b*) on its neck, hollowed

into a spherical cavity at (*c*) to fit the spherical valve of glass (*d*). This valve is set loosely in the brass piece (*e*), that it may, when pressed into the cavity, accommodate itself to its proper place. The brass piece (*e*) has a long tail (*f*), which passes freely through the milled-headed screw (*g*), and is surmounted at (*f*) with a small button, with which the valve may be raised, and which prevents it from falling out, when the whole frame is detached from the collar (*b*), in order to empty or fill the upper vessel (*a*, fig. 1). On the stem (*f*) of the valve (*d*), below the milled screw, a coiled spring wire is fixed, the pressure of which on the valve may be regulated by the milled screw. The flanch of the frame (*h*) has notches cut into it, which fall into corresponding notches in pieces fixed to the ring of the collar. The notches and the whole flanch pass under these pieces with a circular motion, and are thus secured till turned round in such a manner, as that the notches shall again correspond.

When the collar is not secured by the valve, it drops loosely down on the neck of the upper vessel (*a*), as represented in the figure at (*x*).

Fig. 6, (Pl. XIV.) an enlarged view of the hollow glass piston. (*a*) part of the piston rod. (*bb*) two small rings to hold the packing. (*c*) the packing. (*d*) the valve, an unpolished plane convex lens, fitted to the end of the piston. (*f*) the plan of the end of the piston, the latter is placed on the plane side of the lens. (*eeee*) notches in the upper small ring of the piston, over which are passed crossed threads, in order to confine the valve (*f*), and to prevent it in action from being carried off the end of the piston. The notches are necessary to guard the threads from the attrition of the barrel.

Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.



Fig. 6.



Fig. 7.

Fig. 8.



Fig. 8.



Fig. 5.

Fig. 7, (Pl. XIV.) (*a*) a glass vessel with only one aperture to be used instead of the upper vessel (*a*) when the safety valve is considered unnecessary. For the same purpose the receiver (*e*) figs. 1 and 2, may also be used occasionally, as all the joints fit each other. (*b*) a glass stop-cock to be joined occasionally by a pair of collars to the vessel (*a*) in this figure, or to any other at pleasure. This vessel (*a*), with its stop-cock being joined to the bottom of the piston (*d*, fig. 1.), and in the place of the receiver (*e*) may be exhausted if required. But in order to apply the instrument to the purposes of exhaustion, the bottom of the piston should be set in a platform, and have under it a bridge reversed, similar to that represented in fig. 4.—If the instrument were so constructed, bent tubes with rings of glass and collars might be applied by their flat joints, and would answer for exhausting jars connected with them.

Fig. 8, (Pl. XIV.) tubes differently bended, which may be connected at pleasure by flat joints, so as to form syphons of any size, and conducting pipes in any direction, and of any length.

I have to apologize for the minute particularity of this communication; but as I wish artists to understand me perfectly, I have preferred to be rather prolix, than obscure.

I am, dear Sir, with high esteem and respect,

your humble servant,

GILBERT AUSTIN.

XXI. *On the Formation of Fat in the Intestines of living Animals.* By *Sir* Everard Home, *Bart. Presented by the Society for promoting the Knowledge of Animal Chemistry.*

Read March 18, 1813.

THE investigation of the digestive organs of different animals, in which I have been engaged for many years, has led me imperceptibly into an enquiry respecting the particular uses of the lower portion of the intestines in birds and quadrupeds.

The first thing that attracted my notice more particularly to this subject, was finding that in all animals, whose stomachs are made up of a great variety of parts for the purpose of economizing the food, the colon has a greater extent of surface, and the course of the canal is so disposed, that its contents must be a long time in their passage through it. This circumstance led me to believe that the food, after the chyle is formed and separated from it, undergoes in the lower intestines some changes, by which a secondary kind of nourishment is extracted from it.

This opinion was much strengthened, by finding that the colon of the casuary from Java is only one foot long, and each of the cæca which are appendages to it, only six inches long, and a quarter of an inch in diameter, while the African ostrich has the colon forty-five feet, and each of the cæca two feet

nine inches in length, and at the widest part three inches in diameter: besides which, both the colon and cæca have very broad valvulæ conniventes not met with in the casuary from Java. This wonderful difference, for it is more than fifty to one, can only be explained by the luxuriancy of Java being so great, that this bird might destroy its health by over-feeding, had no guard been furnished by nature.

This guard is, the food passing through the intestines with so much facility, and in so short a time, that, however much the bird may eat, only the necessary quantity of nourishment is carried into the constitution; but in the African ostrich, the food is retained in the extensive colon till every thing nutritious is extracted. In all ruminating animals, the colon is of great length, is fixed in its course, which is very intricate, and varies in every different genus; so that we cannot doubt of some particular process being carried on in it.

The process which the contents of the colon undergo, is quite distinct from any thing carried on in the other intestines, since they entirely change their appearance and smell; and there is commonly a valve to prevent any part of them, even the gases evolved, from being carried up into the small intestines.

The peculiar smell of the fæces, which borders so closely on that from putrefaction, although by no means the same, led me to compare them with the animal matter buried in the earth, which is converted into adipocere: in both cases the substance is in the incipient state of putrefaction, but that process never completely takes place; it is excluded from the external air, is either under water, or within the reach of imbibing moisture; and there is no substance whatever, the chyle

excepted, which can better supply the waste produced by the actions of growth and muscular exertion, than animal fat.

The more I canvassed this new opinion, the greater number of circumstances in favour of it occurred to me; one of the strongest of which is, that there is no other mode I am acquainted with, by which animal fat can be formed. To this may be added the curious circumstance of the sleeping animals, which lay in so large a supply of it, in a short time, to serve for their winter's consumption, having a formation of the intestines almost peculiar to themselves, in which there is no valve to distinguish the colon, and no fixed course for that intestine; so that the contents pass along with more facility, and remain a shorter time in the canal, the food being sufficiently plentiful during the summer to compensate for this want of economy, by which the lower intestines receive more abundant supplies for the production of fat. These intestines remain empty during the sleeping season, so that no fat can be formed in that period.—With this very important information, thus procured, in support of my opinion, I have been led to prosecute this inquiry with increased ardour, and shall now bring forward the facts I have been able to ascertain in confirmation of my hypothesis. These I shall detail in the order in which they were acquired, thinking it better to lay before the Society the regular process of the investigation, than to grasp at once at the conclusions, which in the end of it I have felt myself authorized to draw.

I shall therefore begin by stating the circumstances under which adipocere is formed from animal matter, most nearly resembling those in which the contents of the lower intestines in living animals are placed; and this I shall do from facts,

entirely within my own knowledge, the specimens of the adipocere being now in my possession; and afterwards go on by bringing forward proofs that a substance similar to it is formed in the colon.

MARY HOWARD, aged forty-four, died on the 19th of May, 1790, and was buried in a grave ten feet deep at the east end of Shoreditch church-yard, ten feet to the east of the great common sewer, which runs north and south, and has always a current of water in it, the usual level of which is eight feet below the surface of the ground, and two feet above the level of the coffins in the graves. In August, 1811, the body was taken up with some others buried near it, for the purpose of building a vault, and the flesh in all of them was completely converted into adipocere or spermaceti. In STOWE's History of London, this part of Shoreditch is stated to be a morass, and since that time the ground has been raised eight feet. The clerk and the grave-digger observe, that at the full and new moon the water in the sewer rises two feet, and that at those times, there is water found in the graves, which at other times are dry.

The current of water, which passes through the colon, while the loculated lateral parts are full of solid matter, places the solid contents in somewhat similar circumstances to dead bodies in the banks of a common sewer.

The circumstance of ambergris, which contains sixty per cent. of fat, being found, in immense quantities, in the lower intestines of the spermaceti whales, and never higher up than seven feet from the anus, is an undeniable proof of fat being formed in the intestines; and, as the ambergris is only met with in whales out of health, it is most probably collected

there from the absorbents under the influence of disease, not acting so as to take it into the constitution.

Ambergris is found in lumps from fourteen to more than one hundred pounds each; it is not to be distinguished in its appearance from the fæces, but when exposed to the air, it grows hard: a lump has been found in the sea weighing one hundred and eighty-two pounds.*

In the human colon, solid masses of fat are sometimes met with in a diseased state of that canal, and are called *scybala*; these are in all respects similar to ambergris.

Concretions of olive oil and mucus found in the human intestines must be formed in the same way. A case of this kind was communicated to me by our associate Dr. BABINGTON in the following letter:

" My dear Sir,

17, Aldermanbury, Feb. 2, 1813.

" The following are the circumstances relating to the " change produced upon olive oil, by passing through the " stomach and intestines of the elderly person, whose case I " mentioned to you at the last meeting of our Animal Che- " mistry Society. The lady, in question, had for several years " past suffered from severe affections of the stomach, which, " from the attendant symptoms, were considered as occa- " sioned by the irritation of biliary concretions. Many reme- " dies having been resorted to without affording her other " than temporary benefit, she was advised to try the effects " of olive oil, taken to the quantity of two or three ounces at " a time, and to be repeated as circumstances might require. " From this she experienced almost immediate relief, and, on

* Vide Phil. Trans. 1783.

" the subsequent examination of what passed from the bowels,
" globular concretions were uniformly observed, which by
" the persons about her were considered as the gall-stones,
" which had previously been productive of so much distress.
" This lady having occasion some months since to visit her
" friends in town, and a doubt having been suggested by one
" of her medical attendants in the country, as to the nature of
" the concretions in question, I was desirous, from the account
" that I had received, to have an opportunity of determining
" the point for myself; and therefore requested, that if the
" pain should recur, and she should be under the necessity of
" repeating her medicine, that the concretions, which had
" been said always to pass from the bowels in consequence of
" her so doing, might be reserved for my inspection. In a
" few days I was summoned to make my proposed visit, and,
" upon examining the substances collected, I found their ap-
" pearance to be such as I have already described to you,
" namely, that of distinct globules, varying in size from that
" of a large pea to the bulk of a moderate grape, of a cream
" colour, and slightly translucent, of sufficient consistence to
" preserve their form, and to bear being cut by a knife, like
" soft wax, but at the points of their contact disposed to co-
" here. When exposed to heat, they readily melted, and then
" at once exhibited their original oily character. The change,
" which they have since experienced, has taken place in the
" water in which they have been kept.

" I am, dear Sir,

" yours always, very faithfully,

(Signed) WM. BABINGTON."

To Sir Everard Home, Bart.

Our associate, Mr. BRANDE, afterwards examined the substance, and made the following report upon it.

"The globules voided appeared to be composed of the "olive oil combined with mucus: the latter separated during "putrefaction, and the oil was evolved, apparently unaltered. "The relative proportion may be estimated at one-third ani- "mal matter, and two-thirds vegetable oil."

The following case, which was also communicated to me by Dr. BABINGTON, shews that fat is sometimes formed in the intestines, and detected passing off with the fæces.

ELIZABETH RYDER, four years and a half old, had been healthy for six months after her birth, when she became thin, had a sallow complexion, and was liable to jaundice. At a year and a half old, her belly was tumid, and she had great weakness in her back and limbs, for which complaints Dr. BABINGTON was first consulted. At three years old, her mother observed something come from her, as she walked across the room, which, when examined, was found to be fat in a liquid state, which concreted when cold. Ever since that time to the present, she has voided, at intervals of ten or fourteen days, the quantity of from one to three ounces, sometimes pure, at others mixed with fæces; when voided, it has an unusually yellow tinge, and is quite fluid like oil. Her appetite is good, as well as her spirits, and her flesh firm; her belly rather tumid, but not hard: she is subject to occasional griping: her urine natural, and she sleeps well. The specimen on the table was procured under circumstances which precluded all possibility of deception.

These facts, so strongly in favour of the opinion I had taken up, led me to devise in what way it might be put to the test

of experiment. I tried to extract fat from the contents of the colon in different parts of its course, but without success. Failing in this mode of obtaining any decisive conclusions, I was led to believe the cæca of birds more favourable for experiments on this subject, and had those of a duck examined by Mr. W. BRANDE after the bird had been seven days without an evacuation. This confined state of the bowels put the parts nearly under the same circumstances, as if they were in a diseased state. When the cæca were examined, they were found completely distended with fæces of the consistence of soft clay, so that, when the bags were laid open, the contents retained the same form. The intestine, immediately above the cæca, was empty, but the rectum was much distended; its contents were of a softer consistence than those of the cæca.

The following is Mr. BRANDE's report on this subject.

"The contents of the cæca were divided into two portions, "of one drachm each, and comparative experiments were made "with similar quantities of the contents of the rectum.

"*Exp.* 1. One drachm of the contents of the cæcum was "completely immersed in half an ounce of water, and kept "for seven days in a temperature varying from 40° to 60°. "At the end of that time, warm water was poured upon it, "but no appearance of fat could be perceived.

"*Exp.* 2. The same quantity of the contents of the cæcum "was immersed in water containing one-fifteenth part of "nitric acid, and kept under the same circumstances as in the "former experiment. In seven days, warm water poured upon "it separated a portion of oily matter, which concreted when "cold, and appeared to be one-eighth of the whole mass.

"*Exp.* 3. Portions of the contents of the rectum were "treated in the same way, as in experiments 1 and 2. That "in water became putrid very rapidly, and shewed no ap- "pearance of fat. The other, in the diluted nitric acid, was "more dissolved than in Experiment 2. Considerable extri- "cations of gas took place, but there was no appearance of "fat."

From these experiments, we learn that the contents of the cæcum, being confined there for some days, are in a state readily to be converted into fat by nitric acid, while the contents of the rectum are not, probably from being too putrid.

While engaged in this enquiry, I received from Sir JOSEPH BANKS the carcase of a wild swan, which the Hon. Mr. PELHAM had shot in the neighbourhood of the Humber. On examining the cæca, their contents were found to be of a bright green colour. This led me to propose to Mr. BRANDE to ascertain by experiment, whether an admixture of bile had any effect upon the process of converting animal substance into fat. The following experiments were made by Mr. BRANDE upon this subject.

Exp. 1. He took two portions of human muscle of the same size, and digested one of them in human bile, the other in water, both placed in the temperature of 100°.—In the first day the muscle in the bile underwent no change. On the second day it became soft in its texture, and had a slightly fetid smell. On the third day it became more fetid and yellow. On the fourth it had the smell of excrement, was flabby, very offensive, and fatty upon the surface. The portion of muscle, digested in water, had undergone no other change in

the four days, but becoming slightly putrid, and there was no appearance of fat whatever.

Exp. 2. A similar experiment to that with the human bile, was made with a small portion of beef, and ox's bile, and the results were exactly similar.

Exp. 3. The last experiment was repeated in the temperature of 60°. In four days the beef became slightly fetid, and of a yellow colour; in six days it became more fetid, but there was no appearance of fatty matter.

Exp. 4. A portion of beef cut into pieces was digested in ox's bile, at the temperature of 100°. At the end of the fourth day the putrefaction was more advanced than in Experiment 2; the beef was washed and heated upon paper, but no greasy stain was produced.

From these experiments we learn, that the bile has a power of converting animal substance into fat; and that the temperature of 100°, or nearly so, is necessary for that process. We learn also, that this change is produced just as putrefaction is beginning to take place, and if the substance goes rapidly into putrefaction, no fat is formed, and, what is deserving of observation, the peculiar smell belonging to fæces, so different from that of putrid matter, is produced at the time that fat is procured.

Having succeeded in changing animal matter into fat, by adding bile to it out of the body, I was desirous of ascertaining whether this process could be detected going on in the human intestines, and being in attendance upon a gentleman of an advanced age, who had been six days without an evacuation from the bowels, confined to bed by the gout, I did not let slip the opportunity of his having a very costive stool deeply

tinged with bile, to make the experiment. The excrement was put into water, and kept heated for three hours to a temperature of above 100°. When the water was allowed to cool, a film was formed upon the surface, which appeared to be of an oily nature, and Mr. BRANDE ascertained it to be so. The quantity was not great, but quite sufficient to ascertain the fact, and next day the fæces having subsided, the fatty film was much more conspicuous. In the Phil. Trans. for 1673, p. 6093, a case is stated of a person who laboured under an indisposition, attended with sickness and vomiting. In one attack of vomiting, he brought up matter resembling tallow, four pieces of which weighed half an ounce.

This process of forming fat in the lower intestines by means of bile, throws considerable light upon the nourishment derived from clysters, a fact well ascertained, but which could not be explained. It also accounts for the wasting of the body, which so invariably attends upon all complaints of the lower bowels. It accounts, too, for all the varieties in the turns of the colon, which we meet with in so great a degree in different animals. This property of the bile explains likewise the formation of fatty concretions in the gall-bladder, so commonly met with, and which, from these experiments, appear to be produced by the action of the bile on the mucus secreted in the gall-bladder: and it enables us to understand the following effects, which arose from the circumstance of no part of the bile passing into the intestines.

A child was born, at the full time, of the usual size, and lived for several months, but never appeared to increase in size, although it fed heartily, had regular stools, and the food seemed perfectly digested. There was no bile in the stools,

and the skin was of a dark yellowish brown. I saw the child while it was alive, and was struck with its want of growth, and its having no fat under the skin, which made it appear longer than new-born children generally are. Upon examining the body after death, the only mal-formation met with, was there being no gall-bladder, nor any duct leading from the liver into the duodenum.

From what happened in this case, a supply of fat appears necessary for growth, for the child was by no means wasted in its muscles, which it must have been had the constitution not been supplied with nourishment.

Animal fat has, I believe, hitherto been considered as a secretion, although there is no direct evidence in favour of such an opinion. It has nothing in common with the secretions; it is met with in all the interstices of the body; is very often quickly deposited, and in as short a time taken back into the constitution. In these respects it corresponds with the watery fluids, with which the body is supplied.

In a former communication respecting the stomachs of animals, I explained that water was taken up from the stomach by channels yet unknown, and carried into the circulation; from whence it is poured into all the cavities of the body, or thrown out altogether by the kidneys and glands of the skin.

On the present occasion, I hope that I have collected a sufficient body of evidence to prove, that fat is formed in the intestines, and from thence received into the circulation, and deposited in almost every part of the body. When there is a great demand for it, as in youth, for carrying on the growth of the body, it is laid immediately under the skin, or in the

neighbourhood of the abdomen: when not likely to be wanted, as in old age, it is deposited in the interstices of muscles, to make up in bulk for the wasting of the muscular fibres. There appear to be no direct channels by which any superabundance of it can be thrown out of the body, so that when the supply exceeds the consumption, its accumulation becomes a disease, and frequently a very distressing one.

XXII. *On the colouring Matter of the black Bronchial Glands and of the black Spots of the Lungs.* By George Pearson, *M. D. F. R. S.*

Read February 25, 1813.

In the adult human animal the glands designated *bronchial* are generally of a black or dark blue colour. These organs, which are allowed to be lymphatic glands, as is well known, are situated at the root of the lungs externally, within cellular membrane, near the bifurcated trachea; as well as internally on or near the large branches of the bronchi.

At the age of about twenty years the lungs have a mottled, or marbled appearance, from black and dark blue spots, lines, and points disseminated immediately under the transparent pulmonary pleura. As hath been repeatedly observed, the lungs generally become more dark coloured proportionately to their age. Accordingly at upwards of sixty-five or seventy years of life, they often appear almost uniformly black, from the number and congeries, or coalescence of the maculæ, points, and lines just mentioned. Throughout the whole interior substance of the lungs the black spots are seen in a great measure corresponding to the external appearance.

I do not find that any observations and experiments have been made to determine the nature or cause of the black colour above described of the pulmonary organs. It is true, a conjecture has been proposed, that sooty matter taken in with

the air may be the occasion of the colour of the lungs; and that the colour of the glands is occasioned by a peculiar secretion. But the former conjecture has been supposed to be satisfactorily refuted by the absence of the appearance in question among brute animals; as also by its presence in persons who breathe the air of the provinces at a great distance from towns, or places of great consumption of coal; and the latter conjecture is palpably erroneous, because the bronchial glands, of which I am speaking, are not organs of secretion, but of conveyance of lymph.

The course of investigation, in which I have long been engaged, to improve the pathology of pulmonary consumption, led me to some experiments and observations on the subject now stated, which I respectfully submit to the consideration of the Society.

After cutting away the cellular membrane surrounding the black glands, and washing them till the water was no longer coloured, I subjected them to examination.

1. On pressure between the fingers, to burst the investing coat, a black fluid issued which stained the skin, which rendered water black, and which did not alter in colour or apparently dissolve, even at a boiling temperature, either in water or in concentrated muriatic and nitric acids.

2. On breaking down the structure and triturating in a glass mortar a number of these glands with a small proportion of water, a thick black liquid was produced, which was decanted from off many membranous and fibrous masses. But after repeated affusions and trituration, I could not deprive these masses of their black colour, although the water was at last scarcely tinged: it was only by dissolution in caustic potash

lye, or in nitric and muriatic acids, that I could totally separate the black matter from the animal substance to which it seemingly adhered. After repose, a black sediment took place in the waters of elutriation, as well as from the alkaline, and the acid dissolution, which on decantation and evaporation to dryness, afforded the deposit in the state of a black powder.

The texture and proportion of the tingeing matter of the glands was very different in different subjects; whether the lungs, to which they belonged, were in a healthy or diseased condition. In persons of about eighteen to twenty years of age, some of the bronchial glands contained no tingeing black matter at all, but were of a reddish colour; others were streaked, or partially black; and others were quite black, or of a dark blue colour.

3. By boiling the black glands in lye of caustic potash their structure was destroyed, and a turbid black liquid was produced: from which, on standing during several days, a copious black sediment took place; but the liquid still remained black, after remaining at rest a month; much of the tingeing matter continuing suspended. By dilution with water, this matter deposited in a clear liquid.

4. By liquid muriatic acid, of the specific gravity 1,170, the bronchial glands were dissolved at a boiling temperature, affording a turbid black liquor; but on repose, an abundant black deposit took place from a clear yellow liquid, as well as a quantity of the same matter appearing on its surface. On separating this precipitate, and evaporating to dryness, it became a black powder.

5. Nitric acid liquid, of the specific gravity 1,500, most speedily dissolved the substances under examination, affording

a clear yellow liquid with abundance of black matter on its surface. The separation of this coal-like matter is effected most easily with this acid.

It may not be quite superfluous to mention, that from the bronchial glands, which were not at all coloured with black matter, but were merely red, no black matter was separable by the above mentioned acid, and alkaline solvents.

I next examined the black and blue colouring substance of the surface and interior parts of the lungs.

1. A portion of the lungs beautifully marked with a crowd of these coloured areolæ, spots, points, and lines, after being washed till it no longer imparted a bloody tinge to water, still retained these appearances undiminished. On pressure of these coloured parts between the fingers, the skin was sometimes stained of a black colour.

2. On treating the substance of the lungs, abounding in the above described black and blue parts, in the same manner as the bronchial glands just related, with water, with liquid caustic potash, with muriatic and with nitric acids, similar phenomena, with specially the separation of black matter, was observed. The proportion, however, of this black substance was very much smaller to the animal matter dissolved, than in the former cases.

3. To shew that the black matter is of the same kind in the differently figured black parts of the lungs, it may be worth while to relate a particular experiment.

A thin slice of about three inches square of the surface of lungs exhibiting black lines, maculæ, and points under the transparent pleura, was put into a vessel containing three ounces measure of liquid nitric acid. On this it floated, but

all of it was presently dissolved, except the coloured parts, which retained their respective forms till the vessel was shaken: then they were destroyed, and afforded a coating or stratum of black matter on the surface.

Having collected an adequate quantity of the colouring matter in a powdery state, for examination, I performed a number of experiments, but it seems needless to relate more than a few decisive ones.

1. A little of the well exsiccated black powder being sprinkled upon fused nitrate of potash, deflagration took place as with charcoal of wood, or with soot.

2. The same phenomenon occurred with melted oxymuriate, or chlorite of potash, but at a much lower temperature.

3. The deflagration with nitrate of potash, and also with chlorite of potash, was produced, in a suitable glass vessel, to collect the compounded gas; which was received into lime water, and found to be charcoal acid.

4. A little of the black powder was very easily ignited upon a plate of platina, and was speedily burnt off, smelling like burning animal matter, and leaving a minute residue, sometimes of reddish powder and at other times of white.

5. The powder was ignited in a green glass tube closed at one end, and kept red hot from ten to fifteen minutes, the open extremity being slightly stopt with a clay lute to prevent the admission of air. A white vapour with water was discharged, and a minute portion of animal empyreumatic oil was condensed in the upper part of the tube. There remained on cooling, a fine black powder, which in different trials lost $\frac{1}{7}$ to $\frac{1}{4}$; *i. e.* about $\frac{14}{100}$ to $\frac{25}{100}$ of the original weight of the substance.

6. On treating the coaly powder by exposing it to fire, and with the pneumatic apparatus, the products were always charcoal acid, hydro-carbonate gas, and much water, with generally a little empyreumatic oil, and sometimes a trace of prussic acid, leaving a residue varying in weight as just stated.

From the properties above manifested, I conceive I am entitled to declare the black matter, obtained from the bronchial glands, and from the lungs, to be animal charcoal in the uncombined state; *i. e.* not existing as a constituent ingredient of organized animal solids or fluids.

I mean by the term animal charcoal, what is popularly understood. Of course, I do not mean pure charcoal. Such a state of this substance cannot here be reasonably expected, either from a consideration of the state of it, as inspired from the atmosphere, or from its necessary impregnation with animal matter during its long residence in the lungs. I imagine, no person would hesitate to consider such a coaly substance as the present to be charcoal, if derived from other sources besides the animal economy; it being, as shewn by the preceding experiments, a black, tasteless, infusible powder, indissoluble in muriatic acid, nitric acids, and perhaps all common acids, except the sulphuric; affording as large a proportion of charcoal acid as animal and vegetable charcoal which has been exsiccated at the same temperature, and equally resisting fire in close vessels.

For the purposes of physiology, a few theoretical remarks may, perhaps, be useful. I think the charcoal, in the pulmonary organs, is introduced with the air in breathing. In the air, it is suspended in invisibly small particles, derived from the burning of coal, wood, and other inflammable materials in

common life. It is admitted, that the oxygen of atmospherical air passes through the pulmonary air vesicles, or cells, into the system of blood vessels; and it is not improbable, that through the same channel various matters contained in the air may be introduced. But it is highly reasonable to suppose, that the particles of charcoal should be retained in the minutest ramifications of the air tubes, or even in the air vesicles, under various circumstances, to produce the coloured appearances on the surface, and in the substance of the lungs, as above described. It must also be considered, that innumerable absorbent lymphatic vessels take their rise in the bronchial tubes: for the lungs are more richly stored with lymphatic vessels than any other organ, excepting the liver. When I compared the black lines and black net-like figures, many of them pentagonal, on the surface of the lungs, with the plates of the lymphatic vessels by CRUICKSHANK, MASCAGNI, and FYFFE, I found an exact resemblance. And when I found that these vessels contained charcoal, I judged that it was fair to infer, that the lymphatics of the lungs absorb a variety of very different substances, and especially this coaly matter, which they convey to the bronchial glands, and thus render them of a black, or dark blue colour. Hereafter, among other enquiries, the colour of the large trunks of the lymphatic vessels, just *before* they enter the bronchial glands, and just as they pass out of them, ought to be observed. Also the effect of the charcoal thus conveyed into the thoracic duct, or directly into the blood by the lymphatics from the black glands, is, I presume, worthy of attention.

According to this theory, we can account reasonably for the absence of the black colour of the bronchial glands, and of the

lungs in infants, in children, and even frequently in persons of eighteen or twenty years of age. For the same reason these appearances are seldom observed in any brute animal which has fallen under my observation. On a subject so novel, or at least so much neglected, as that on which I am writing, many facts are wanting to establish, demonstratively, any theory which can be proposed; but I know of none, at this time, which are at variance with that I have ventured to offer.

George-street, Hanover-square, June 13, 1813.

P. S. Since my last communication to the Society, I have had an opportunity of making some further observations and experiments on this subject.

The lungs of an infant, who lived two days, were obligingly sent to me by my friend Mr. C. M. CLARKE. I found the bronchial glands quite white, and the lungs appeared externally of an uniform reddish colour. The pulmonary organs of a girl of fifteen years old, I found rather thickly mottled, but the bronchial glands were only tinged on their surface, and did not yield one-fourth of a grain of coaly powder. In two other females also, aged nearly fifteen years, who died of pulmonary consumption, the lungs were not at all marbled, but some of the bronchial glands were tinged black, and others were white. The lungs of two men, the victims of pulmonary phthisis, at the age of twenty-one years, exhibited thinly, black spots, streaks, and areolæ, with many of the glands in question of a deep blue colour. A woman's lungs, thirty-one years of age, were found studded beautifully with black spots, and lines; the bronchial glands being all either blackish

or blue. In no instance have I observed the lungs and glands, here spoken of, so black, and from which I separated so much charcoal, as in those of a person forty-two years old, whose death was occasioned by most extensively diffused tubercles, many vomicæ, and a considerable condensation of the pulmonary organs. I now recollect, that this subject had been a Smoker of tobacco, generally several times, but always once a day, for perhaps more than twenty years. Future observations must determine more satisfactorily the state of the pulmonary organs, according to the impregnation of the air with sooty vapours. If, hereafter, it be shewn, that the lungs of persons living remote from sources of such vapours, are still greatly impregnated with coaly matter, the just conclusion can only be, that such matter is more extensively diffused through the atmosphere, than is apprehended. This being the fact, it would also afford a proof that it is only the invisibly small particles which are absorbed, for the larger particles remain unabsorbed, entangled in the mucus lining the air vessels, and never get farther, but are rejected from time to time by expectoration. Accordingly, in a morning, healthy people, after the night's rest, very commonly hawk up mucous matter of a bluish colour with black streaks, owing to charcoal; and persons in a diseased state, especially by great exertions in coughing, frequently expectorate matter spotted and streaked with black particles. The quantity of coaly matter in the pulmonary organs is not entirely according to the age, for I was disappointed, on finding the lungs and glands in a woman of seventy-five years of age, in London, not more deeply coloured, than is usual at the age of fifty. At present, I am unable to state any connection between certain diseases, and the presence of coaly matter.

Farther investigation has shewn, that this coaly matter does exist in domesticated brute animals; but as they die, or are killed generally before they attain to twenty years, or even fifteen years of age, the organs in question are seldom seen blackened. However, in diseased conditions, the cases are not according to this rule. With the approbation of Professor COLMAN and Mr. SEWEL, several of the worthy students of the Veterinary College have frequently obliged me by furnishing, for my examination, the lungs of horses and asses. In general, the bronchial glands were merely white, or reddish; but now and then they were partially black. In one instance of an ass, only six months old, these glands were black from coaly matter; but the lungs were uniformly red. The animal had died of peripneumony. In no instance have I seen, in any brute creature, the lungs marbled and streaked with black lines as in the human. There is seldom an opportunity of inspecting horses which die from their natural age, viz. of thirty to forty years: for I am informed they mostly die or are killed in London before they are fifteen or sixteen years old. I have not seen coaly matter in the lungs, or glands of the ox kind, sheep, and hogs. The black appearances produced by distended blood-vessels and by ecchymosis, should be recollected, to avoid the error of ascribing them to charcoal. The absence of this matter in human creatures, at the ages just mentioned, when animals are slaughtered or die, affords a proof, although not a decisive one, that the exemption is more reasonably ascribable to the circumstance of time, and living in the open air, than to the peculiarity of the economy of each species of live being. Consistently with this observation, in the instance of a cat, known to have lived in

Mr. THOMAS's family, at least eighteen years, the bronchial glands were quite black from coaly matter, and the lungs were uniformly red; but in all other examinations of much younger cats, I found these glands either white or red.

The blackness of the lungs from charcoal remains, although hæmorrhage to occasion death has occurred. It is not removable by ablution, or maceration in water, nor by acids, nor alkalies, nor by the early stages of putrefaction. I have not met with a similar coaly substance in any parts of the animal economy, except the lungs. The glands of the meso-colon are sometimes black, similarly to the bronchial; but the colour soon disappears on immersion in nitric or muriatic acids, no charcoal being separable. The black, or more truly the dark brown tingeing liquid of the sepia, I have ascertained by experiments, does not contain uncombined charcoal; this matter existing there only as a constituent ingredient of animal matter.

As I have represented, it is conceived that the coaly matter is very slowly absorbed by the mouths of the lymphatic vessels in the innumerable air tubes and cells.

To determine whether or not this matter exists in these lymphatic vessels, and is the occasion of the black maculæ, streaks, and areolæ, or marbled appearance of the surface of the lungs, I entreated Mr. WHARRIE, of St. George's Hospital, whom I knew to be a skilful anatomist, to inject these vessels with quicksilver. In some trials, the injection passed without interruption, in the usual manner; but in others it was apparently obstructed, by meeting with the black lines on the surface. Mr. GEORGE EWBANK also, at my desire, very dexterously dissected out about one inch in length of one of these

black lines, supposed to be a lymphatic vessel. Being put into a glass capsule, full of nitric acid, the black line immediately was contracted in all dimensions; but it retained its form after digestion, for several days, at a high temperature: afterwards on gently shaking the capsule, the black line was broken into a number of indissoluble particles. In the interior of the lungs, it is not unusual to see black spots in the middle of tubercles, although these substances consist apparently of self-coagulated lymph probably secreted in the cellular substance, and therefore very likely to envelope the coaly matter in the air tubes by the coalescence of numerous minute tubercles.

It has been objected, that the nitric acid may develope the constituent or combined ingredient charcoal of all animal substances; and consequently no proof will be thereby afforded of this matter being extraneous; but on many trials, I have never by this means obtained charcoal from any animal mucilage.

I have no reason to believe, that any of the coaly matter under investigation, is dissolved by this acid, for on distilling a pint measure of it, from ten grains of the black powder of the bronchial glands, there was no sensible diminution of it, whether it was so treated before ignition, or subsequently: on evaporation to dryness, nitric acid, in which the coaly matter had been boiled, afforded no black sediment. Hence, I conceive, that this menstruum may be employed to determine, more accurately and speedily, the proportion of the coaly matter, than any other agent. Sulphuric acid does dissolve a certain portion of this charcoal, affording a transparent liquid, even on dilution with water.

XXIII. *Experiments on the Alcohol of Sulphur, or Sulphuret of Carbon.* By J. Berzelius, *M. D. F. R. S. Professor of Chemistry at Stockholm; and* Alexander Marcet, *M. D. F. R. S. one of the Physicians to Guy's Hospital.*

Read May 13, 1813.

THERE has been, of late years, much discussion respecting the nature of a singular oily liquid, which was first noticed by Mr. LAMPADIUS,* who procured it by distillation from a mixture of pyrites and charcoal, and gave it the name of *alcohol of sulphur*, on account of its very great volatility.

LAMPADIUS considered this liquid as a compound of sulphur and hydrogen; but Messrs. CLEMENT and DESORMES,† who obtained the same substance by subliming sulphur through red hot charcoal, were led by their researches to conclude, that the alcohol of sulphur was a combination of sulphur and charcoal, and that hydrogen was not one of its constituent principles.

Doubts, however, were entertained respecting the chemical nature of this compound. Mr. BERTHOLLET believed it to be a triple combination of sulphur, charcoal, and hydrogen.‡ Messrs. VAUQUELIN and ROBIQUET,§ from a joint enquiry on the subject, considered it as a binary compound of sulphur and

* CRELL'S Annals, 1796. II.

† Annales de Chimie, An. X. Vol. XLII. p. 121.

‡ Ibid. p. 296.

§ Ibid. Vol. LXI. p. 145.

hydrogen; and the late Mr. BERTHOLLET, jun.* who resumed the subject, with much apparent accuracy, was induced, from his experiments, to adopt the last mentioned opinion, that the alcohol of sulphur was a compound of sulphur and hydrogen, and that no carbon entered into its formation.

Sir HUMPHRY DAVY, in the course of his brilliant career of chemical discovery, has repeatedly noticed this singular substance;† but it does not appear that he ever examined it minutely, and if he gave the preference to the opinion of the younger Mr. BERTHOLLET, respecting its chemical nature, it was principally from the circumstance of its having apparently yielded sulphuretted hydrogen by the agency of the Voltaic electricity, and sulphureous and sulphuric acid by combustion in oxygen.

These varieties of opinions, amongst such respectable authorities, having left on our minds much doubt and uncertainty with regard to the real nature of this compound, we undertook to examine it conjointly; in hopes that we might succeed in ascertaining its composition with a greater degree of certainty than our predecessors had done, and perhaps also in discovering the exact proportions of the elements of which it is composed, a circumstance which, in the present state of chemical science, would add considerable interest to an enquiry of this kind.‡

* Mémoires d'Arcueil, Vol. I.; and Annales de Chimie, Vol. LXI. p. 127.

† Phil. Trans. 1809, page 464; and Elements of Chemical Philosophy, page 283 and 510.

‡ This enquiry was undertaken in London, in the months of July, August, and September, during Mr. BERZELIUS's stay in this country; and the leading points of the analysis were then ascertained, and mentioned to our chemical friends. Some parts of the work, however, and in particular the precise determination of proportions, remained to be completed; and this has been accomplished by Professor BERZELIUS

Whilst we were finishing this paper, and arranging our materials, in order to present them to the Royal Society, the Annales de Chimie, for the month of September last, reached this country, by which it appears that the alcohol of sulphur has recently been again submitted to examination in France, by Mr. CLUZEL, who supposed it to be a compound of sulphur, carbon, hydrogen, and azote.* But Messrs. BERTHOLLET, THENARD, and VAUQUELIN, the reporters of these experiments, have repeated them, and by a process of their own, quite different from the methods which we have employed, these celebrated chemists have revived the opinion of CLEMENT and DESORMES, that the alcohol of sulphur is a compound of sulphur and carbon, in the proportion of about eighty-five parts of the former to fifteen parts of the latter, no hydrogen entering into its composition. How near this conclusion agrees with our own, will be seen in the following pages.

Although some of the experiments on this substance which we are going to relate, have already been performed by others, yet as we have, in several instances, obtained different results, and as any considerable omission would interrupt the thread of our narrative, we shall beg leave to lay before the Society a complete, though concise, account of our analysis.

We shall, for the sake of arrangement, class the particulars of our enquiry under four heads: viz. 1. *Preparation and general*

since his return to Stockholm. There are also some important collateral objects, which are almost entirely his own, and which, together with his remarks on determinate proportions, have been added to the paper in the form of an appendix. (Note of Dr. MARCET.)

* It appears that Mr. CLUZEL considered the sulphur in this oil, as being in a deoxydated state, an idea which had some years ago occurred to one of us, and also to Sir H. DAVY. See Phil. Trans. for 1809, p. 465.

properties of the Alcohol of Sulphur. 2. *Experiments to ascertain whether Hydrogen be present in the Alcohol of Sulphur.* 3. *Experiments to ascertain the presence of Carbon.* 4. *Experiments to ascertain the proportions of the elements of the Alcohol of Sulphur.* And the Appendix will contain various elucidations and collateral objects.

§ 1. *Preparation and general Properties of the Alcohol of Sulphur.*

We prepared this substance according to the method pointed out by CLEMENT and DESORMES, that is, by slowly volatilizing sulphur through red hot charcoal in a porcelain tube, and condensing in water the oily liquid which is thus formed. The details of the process have so often been described elsewhere, that it would be superfluous to repeat them here. The fluid thus procured is of a pale yellow colour; being extremely volatile, it produces a remarkable degree of cold during its evaporation, and deposits in a crystalline form, some sulphur which it held in solution. The quantity of sulphur thus dissolved in the oily liquor, varies according to the circumstances of the process by which it is obtained;* and its specific gravity, before it has been rectified, must, of course, be liable to corresponding variations.†

In order to obtain the alcohol of sulphur perfectly pure, it is only necessary to distill it at a very gentle heat, not exceeding 100° or 110°; and some dry muriate of lime may be put into the retort, in order to obtain the liquor perfectly free from moisture. The fluid which comes over is quite pure, and some sulphur remains in the retort.

* The alcohol of sulphur can dissolve as much as ¼ of its weight of sulphur.

† We ascertained the specific gravity of only one specimen of the impure oily liquor, which proved to be 1,321.

The alcohol of sulphur thus prepared has the following properties: it is eminently transparent, and perfectly colourless. Sometimes, immediately after distillation, the oily liquid appears somewhat opaque and milky; but the next day the liquor is found perfectly clear, the milky appearance having spontaneously disappeared. It has an acrid, pungent, and somewhat aromatic taste; its smell is nauseous and fetid, though distinctly differing from that of sulphuretted hydrogen. Its specific weight is 1,272; its refractive power, as ascertained by Dr. WOLLASTON, is 1,645. Its expansive force (the height of the barometer being thirty inches, and the temperature 53.5° FAHRENHEIT) is equal to the pressure of 7,96 inches of mercury; so that air, to which it is admitted, will dilate about one-fourth of its volume. It boils briskly under the common atmospheric pressure at a temperature between 105° and 110°.* It does not congeal at a temperature as low as 60° below zero of FAHRENHEIT's scale. It is highly inflammable, and takes fire at a temperature scarcely exceeding that at which mercury boils; it burns with a bluish flame, emitting copious fumes of sulphureous acid. If a long glass tube, open at both ends, be held over the flame, care being taken to keep the tube quite cold, no moisture whatever is deposited on its internal surface.

This oily liquid readily dissolves in alcohol and ether,† though not in all proportions; and if it contain sulphur in solution, the

* The volatility of this liquid is very remarkable; it exceeds considerably that of ether; and in some experiments, tried since this paper was presented to the Society, on its power of producing cold by evaporation, some unexpected results have been obtained, which may, perhaps, become the subject of some future communication.

† Ether can dissolve about three times its own bulk of the oily liquor before any separation or turbidness takes place.

latter is rapidly precipitated in spicular crystals. The spirituous solution is rendered turbid, and the oily liquor precipitated by the addition of water. The alcohol of sulphur is not soluble in water, though it imparts to it some of its odour; this appears to be owing to its impregnating the atmospheric air contained in the water, rather than the water itself. It readily mixes, and perfectly incorporates with either the fixed or volatile oils; but if it hold sulphur in solution, the latter separates in a crystalline form. It dissolves camphor very rapidly, and forms with it perfectly transparent solutions. When kept for some time under water, the latter being in contact with atmospheric air, neither the air, nor the oily liquid, suffer any alteration. On being heated in contact with potassium, the alcohol of sulphur suffers no change or decomposition, when in its liquid state; but if it be brought to the state of vapour, by the exclusion of the air, and heat applied, the potassium becomes ignited in the vapour, and emits a reddish flame. During this species of combustion, a blackish film appears upon the potassium, (as happens when potassium is burnt in carbonic acid gas), and on introducing water, a greenish solution is obtained, which contains a quantity of carbonaceous matter, and has the smell and other properties of hepar sulphuris. Neither mercury nor amalgams of silver, or of lead, are at all acted upon by this liquor, if it has been carefully distilled; but if it contain any sulphur in solution, the amalgam blackens, and sulphurets of silver, or of lead, are produced, after which the liquor remains pure, and unsusceptible of further alteration. Mercury heated to ebullition in an atmosphere of this oil, comes over unaltered, and without the liquor undergoing any change. The alkalis dissolve it entirely, though but very slowly. None of the

acids exert any sensible action upon this liquor, except the nitro-muriatic acid, and oxy-muriatic gas in a humid state. If a globule of the alcohol of sulphur be brought into contact, under water, with a globule of the oily fulminating compound, lately described to this Society by Sir H. Davy, the two globules remain in contact, side by side, without mixing, and without exerting any action upon each other; but when stirred together, they incorporate, forming a homogeneous amber-coloured globule, which does not detonate, even when exposed to ignition; and if olive oil be brought into contact with the two oils thus mixed, no detonation or other obvious effect takes place, a remarkable circumstance, of which advantage might be taken to attempt the analysis of the fulminating substance. The detonation of the fulminating compound with phosphorus, is also prevented by the presence of alcohol of sulphur; but if the quantity of the detonating compound preponderates over that of either the olive oil or phosphorus, the mixture inflames, though without detonation.*

§ 2. *Experiments to ascertain whether Hydrogen be present in the Alcohol of Sulphur.*

1. We introduced into Volta's eudiometrical tube, over mercury, some pure oxygen gas, with one drop of the oily liquor. After letting these stand together for a few minutes, an electrical charge was passed through the tube, which produced a vivid explosion. The gas was first reduced to between one-

* These experiments were tried in the presence of one of us, by Mr. Wilson, Assistant to the Chemical Lectures of Guy's Hospital, who, conjointly with two other gentlemen, has lately published in Nicholson's Journal, many curious facts on this extraordinary compound.

fifth and one-sixth of its volume, a circumstance which appeared to be owing to the destruction of the volatilized portion of the liquor; for the gas soon afterwards recovered its former volume. On examining the superior part of the tube, we perceived traces of a condensed liquid. Whether this liquid was concentrated sulphuric acid, or simply a small portion of the oily liquor, remained to be determined. We repeated the experiment in the following manner.

A quantity of the oily liquor was suffered to evaporate in oxygen gas of known purity, and previously dried with muriate of lime. Some of this gas was introduced into the eudiometer, and exploded by the electrical spark. It lost between one-fifth and one-sixth of its volume; but we perceived, even then, traces of a condensed liquid in the superior part of the eudiometer. This liquid, in about a quarter of an hour, became opaque, and ultimately formed white specks, which we found to be sulphate of mercury. No oxygen gas remained in the tube after the explosion; but the gaseous residue consisted of sulphureous acid gas, and, as we shall see afterwards, of the carbonic acid, and the carbonic oxyd gases. This experiment having still left undetermined, whether the vestige of liquid we had observed, was or was not to be ascribed to the formation of water, we tried to decide this point by means of oxymuriatic gas, in the following manner.*

2. We caused a succession of bubbles of oxymuriatic gas,

* In relating the experiments in which oxymuriatic gas (or chlorine) is concerned, we have used the old nomenclature, and have explained the phenomena according to the notions, respecting the nature of this agent, which prevailed previous to Sir H. DAVY's ingenious experiments and speculations on the subject. But those who consider his views, as sufficiently established to supersede entirely the old hypothesis, may easily adapt to our statements the language which belongs to the new doctrine.

previously dried by muriate of lime, to pass through a portion of the alcohol of sulphur, and afterwards through a quantity of distilled water, over which it was collected. The oily liquid suffered no other change than that of acquiring an orange hue. After an hour and a half the process was stopped, and the liquor was found to have absorbed a considerable quantity of the gas, which had imparted to it a peculiar and extremely strong odour. Most of the gas, however, had passed through the water, in which it had deposited a portion of the oily liquid unaltered. The water had acquired a peculiar smell, and contained, after the expulsion of the oxymuriatic gas, a little muriatic acid, with a vestige of sulphuric acid. Here again it remained doubtful, whether the production of muriatic acid was owing to the alcohol of sulphur containing hydrogen, which, by uniting with the oxygen of the oxymuriatic acid, had formed water; or whether this appearance of minute quantities of muriatic and sulphuric acid, might not be explained in some other manner. It will hereafter be seen how the last conjecture was verified; but in the mean time it was sufficiently shewn, by the experiment just related, that if the alcohol of sulphur really contain hydrogen, it must be in very small quantity, and probably from some accidental circumstance, since the greater part of the liquor remained undecompounded, and with no other alteration than the absorption of the oxymuriatic gas. In the course of a few days, however, the oil gradually lost the smell of this gas, and acquired that of the muriated sulphur, described by Dr. Thomson, (Sir H. Davy's sulphurane). The liquor thus treated, on being exposed to the action of water, lost its colour, and resumed its original

characters; and this was accomplished more quickly, if the water contained some alkaline substance.

3. Into a glass receiver, full of oxymuriatic gas, we immersed some of the oily liquor previously ignited in the air; it was instantly extinguished, and appeared to undergo no change, except that of absorbing a little oxymuriatic acid, and thereby acquiring a yellow colour.

4. A portion of the oily liquor, in the state of vapour, was caused to pass through liquified muriate of silver heated to a cherry red, and to condense again into a small receiver artificially cooled; neither the liquor, nor the muriate of silver, were altered by that operation, nor did the air contained in the apparatus appear, on examination, to contain the smallest quantity of acid. This shewed that the liquor did not contain any hydrogen, since if it had, the muriate of silver would have been decomposed, the hydrogen uniting with the oxygen to form water, so as to generate muriatic acid gas (which may be considered as a muriate of water, or more correctly a muriate of hydrogen), whilst some sulphuret of silver would have been produced.

5. Though the result just related appeared sufficiently conclusive, we thought it desirable to have it confirmed by some other mode of proceeding. With this view, we heated to incipient redness, in glass tubes, various metallic oxyds, such as red oxyd of iron, black oxyd of manganese, oxyd of tin; and we caused quantities of the oily liquor in vapour to pass through these ignited oxyds. The liquor was, by that means, entirely decomposed; the metallic oxyds were converted into sulphurets, and the gases acquired a strong smell of sulphureous

acid. But we could not, in any of our experiments, detect the least production of water, although our apparatus was so devised, as to render the smallest quantity of water conspicuous, and although we burnt, in some instances, as much as fifty or sixty grains of the liquor.

From all these experiments, we think ourselves warranted in concluding, that *the alcohol of sulphur contains no hydrogen.*

§ III. *Experiments to ascertain the presence of Carbon in the Alcohol of Sulphur.*

1. The gaseous residue obtained from the combustion of the vapour of the alcohol of sulphur in oxygen gas, by means of VOLTA's eudiometer (§ II. 1.), being put in contact with water, was in a great degree absorbed, and the water acquired the taste and smell of sulphureous acid. The remaining gas being agitated with lime water was partly absorbed, and produced a precipitate of carbonate of lime. The unabsorbed portion being mixed with oxygen gas, and the electrical spark passed through the mixed gases, they detonated, and the remaining elastic fluid was found to have again acquired by this detonation the property of rendering lime water turbid, and of forming carbonate of lime. The residue of the first detonation was therefore gaseous oxyd of carbon.

2. We introduced into a glass receiver, filled with pure oxygen gas, and inverted over mercury, a small glass capsule full of the oily substance, which we kindled in the air before we plunged it into the gas. It continued to burn, and we were not a little surprised to find, that the heat of this combustion was sufficiently intense to melt a pretty strong platina wire, by which the capsule was suspended, so that it fell with its

contents upon the surface of the mercury, where the oily liquor continued to burn until it was all consumed. Water being then introduced, an absorption of gas took place, and the water became sulphureous. The unabsorbed portion of gas being introduced into lime water, rendered it turbid, and formed a quantity of carbonate of lime, which was sufficient to be carefully examined and accurately estimated.

3. We caused some alcohol of sulphur to dissolve in barytic water, in a well stoppered bottle. This solution proceeded very slowly, and was only completed at the end of three weeks. The barytic water had assumed a yellowish colour, and had deposited a white precipitate, which being well washed, and treated with liquid sulphureous acid, dissolved in it with effervescence. It was therefore carbonate of barytes.

4. A similar experiment being tried with lime water, an analogous result was obtained.

It follows from these experiments, that *the alcohol of sulphur contains carbon*, and therefore that this body, the nature of which has been the subject of so much doubt and speculation, is a true SULPHURET OF CARBON, a name by which, in compliance to the received chemical nomenclature, we shall henceforth usually designate it in the course of this paper.

§ IV. *Experiments to determine the Proportions of Sulphur and Carbon, in the Sulphuret of Carbon.*

In attempting to analyse, with accuracy, the sulphuret of carbon, considerable difficulties occur from the great volatility of that compound, and from the inconsiderable action which most chemical agents exert upon it.

1. We first tried to ascertain the proportion of its constitu-

ent parts, by exploding it, in a state of vapour, with oxygen gas, in Volta's eudiometer; but this method presented insurmountable obstacles, and in particular that of our not being able to determine, with accuracy, the quantity of the oily substance actually decomposed by that process.

2. On the other hand, on attempting simply to burn in oxygen gas the sulphuret of carbon in its liquid state, the necessity of previously kindling it in the air, in order to avoid explosions, prevented our being able to ascertain, with accuracy, the quantity of the oily substance submitted to the experiment.

3. We also attempted the oxydation of the sulphuret of carbon by nitro-muriatic acid; but we obtained by this method, as we shall see afterwards, results and combinations which were quite different from those which we had expected.*

4. We then had recourse to the action of alkaline substances on the sulphuret of carbon, with a view to produce an alkaline sulphuret, and afterwards to convert this into a sulphate by means of nitro-muriatic acid. The sulphuric acid, precipitated by a barytic salt, was ultimately expected to indicate the quantity of sulphur contained in the oily compound. In order to try this method, we exposed portions of the sulphuret of carbon to the action of caustic potash, ammonia, barytic water, &c. and when the solutions appeared to be completed, we tried the effect of acids upon them; these separated from them a soft, reddish, glutinous substance, which, when exposed to distillation, yielded, first, a portion of the oily liquor undecomposed, and afterwards sulphur; thus shewing that the

* A new and singular compound was obtained from the long continued action of nitro-muriatic acid on the alcohol of sulphur, the history and composition of which will be detailed in the Appendix.

whole of the sulphuret of carbon had not been decomposed. A black powder remained in the retort, which was found to be charcoal, and afforded us a new and unequivocal proof of the presence of carbon in the oily substance.

This imperfect decomposition of the sulphuret of carbon in alkaline solutions, was found an insurmountable objection to this mode of analysis; and induced us to try the method of distilling the sulphuret through ignited metallic oxyds.

5. For this purpose, we introduced a glass tube (BC), in an inclined direction, through a small stove (fig. 1, A), the

Fig 1.



Fig. 2.



tube being protected by lute, and filled with red oxyd of iron coarsely pulverized. The sulphuret of carbon, after being accurately weighed, was enclosed in a little retort (E), which we hermetically sealed to the upper extremity of the tube (B). To the other extremity (C), we adapted another glass tube

(CGI), terminating in a mercurial pneumatic apparatus (H), and having a bulb or expansion (in G), at about half way between its extremities, which bulb was cooled down to about 15° FAHRENHEIT, and kept at that temperature during the experiment by means of a mixture of ice and salt. The tube in the stove (BC) being now heated to redness, and an extremely gentle degree of heat applied to the retort, the oily liquor distilled slowly through the red hot oxyd of iron, and the whole of the experiment succeeded in a most satisfactory manner; for we found that the joints of the apparatus had remained perfectly tight during the process, and every vestige of the sulphuret of carbon had been decomposed.* We could not perceive in the bulb the least appearance of moisture.

6. The oxyd of iron (the whole weight of which was about ten times that of the oil decomposed), was partly converted into sulphuret of iron, having a yellow colour and a brilliant metallic lustre. This sulphuret was dissolved by nitro-muriatic acid, and the oxyd of iron was separated by ammonia. The filtered liquid, being neutralized by muriatic acid, was precipitated by muriate of barytes, and the precipitate was well washed, and heated to redness.

7. During the process of decomposition above related, the mixture of sulphureous acid gas and carbonic acid gas had collected in the receiver (H) placed on the extremity of the

* The process was so slowly conducted, that it required six hours and a half for the decomposition of about fifteen grains of the oil. The heat applied to the retort was merely that which radiated from a small lamp (F), placed at some distance from it, a screen being interposed between the furnace and the retort. In several previous trials, we had failed from not being sufficiently cautious and gradual in the mode of warming the retort.

tube. In order to separate these two gases from each other, we made use of the brown oxyd of lead, a substance which one of us had often employed for the purpose of combining with, and separating from other bodies, liquid sulphureous acid, with which that oxyd forms a neutral sulphate. The same method, to our great satisfaction, succeeded perfectly with the gaseous sulphureous acid. After a contact of about one hour, between the brown oxyd of lead and our mixed gases, the whole of the sulphureous acid gas was so completely absorbed, that the carbonic acid gas which remained in the vessel, had not the least sulphureous smell, and yet the carbonic acid itself remained perfectly unaltered in its bulk, after continuing in contact with the oxyd of lead for the space of three days.*

8. The mode in which the respective weight of the two acids was ascertained was this; the oxyd of lead was placed in a small glass capsule (*a*, fig. 2), the orifice of which was carefully secured, by means of a thin piece of glove-leather, against the admission of quicksilver. A very thin flexible metallic wire was fastened to the capsule, so as to enable the operator to raise it through the mercury in the receiver containing the gases, as is expressed in the figure (fig. 2). After continuing the contact for several hours, and when the absorption had

* The brown oxyd of lead, used in our experiments, was recently prepared, and had been long digested in weak nitric acid. In wiping this oxyd, and pressing the moisture out of it, we found it expedient to reduce it to the state of small laminæ, as in this form it is more easily penetrated by the acid gas than in the state of powder. It may be proper to remark, that when, for the sake of trial, gaseous mixtures, consisting of sulphureous acid gas and sulphuret of carbon in vapour, were used, the latter, as the absorption of the acid gas proceeded, was deposited in its liquid state upon the oxyd of lead.

ceased for a considerable time, the capsule was withdrawn, and another similar capsule filled with dry caustic potash (hydrate of potash), was introduced in a similar manner. The remaining gas (with the exception of a small residue),* was immediately absorbed. Both capsules, after wiping off minute particles of quicksilver which adhered to their surface, were carefully weighed and were found to have undergone an increase of weight, which, of course, expressed the quantity of acid they had respectively absorbed. By combining these results with the examination above described (§ IV. 6.), of the sulphuret of iron found in the tube, the proportions of sulphur and carbon were deduced. Two such experiments were made which yielded very similar results; and by a mode of computation, which will be detailed in the Appendix, we were led to conclude that the alcohol of sulphur, or sulphuret of carbon consists of

Sulphur	-	84,89	or 100,00
Carbon	- -	15,17	17,89.

If these proportions be compared with the numbers 13 and 5, which are assumed by Mr. Dalton as representing the respective weights of a particle of sulphur and a particle of carbon; or with those of 30 and 11,4, which are considered by Sir Humphry Davy as representing the same substances, we shall find that either of these modes of computation, as well as that detailed in the Appendix, make the above-mentioned proportions of sulphur and carbon, in the alcohol of sulphur, correspond very nearly to two atoms or portions of sulphur, to one of carbon; a result which is perfectly agreeable to the doctrine of determinate proportions. And as the

* This small residue corresponded almost exactly to the quantity of air expelled from the tube in which the decomposition had been performed.

sulphur and carbon recovered by our analysis, were precisely equal in weight to the sulphuret of carbon subjected to examination, we are the more confident in believing that it does not contain any other element.

APPENDIX BY PROFESSOR BERZELIUS.

A. *Particulars respecting the Mode in which the Proportions of Sulphur and Carbon, in the Alcohol of Sulphur, were established.*

Two analytical experiments were made in succession, in the manner described in the paper (§ IV.), with the following results.

Exp. 1. 1,05 parts * of sulphuret of carbon produced 0,24 parts of sulphureous acid gas, and 0,59 parts of carbonic acid. The sulphuret of iron found in the tube was dissolved in nitro-muriatic acid; the oxyd of iron was precipitated from this solution by caustic ammonia in excess, and the filtered liquor was neutralized by muriatic acid. From this solution, muriate of barytes precipitated a quantity of sulphate of barytes, which, after ignition, weighed 5,6. From Mr. T. DE SAUSSURE's experiments, as well as from inference from the doctrine of determinate proportions, 100 parts of carbonic acid contain from 27 to 27,1 parts of carbon. Therefore the above mentioned 0,59 parts of carbonic acid contain 0,1593 of carbon. From the experiments on the composition of sulphate of barytes, published by myself, 100 parts of that salt contain 13,66 of

* The weights represented by these numbers were *grammes*; so that the quantity of sulphuret of carbon employed in this experiment, was 1,05 gramme, corresponding to about 15¾ English grains.

sulphur; therefore the 5,6 parts of sulphate of barytes, are equivalent to 0,765 of sulphur, which, added to the 0,1224 parts of sulphur contained in the 0,24 parts of sulphureous acid gas, make 88,74 parts of sulphur. The sum total of sulphur and carbon is, therefore, 88,74 + 15,98 = 104,67. But the quantity of the sulphur of carbon was 105 parts; consequently there is a loss of $\frac{1}{3}$ per cent. From this experiment therefore, the sulphuret of carbon is composed of 15,7 parts of carbon to 84,83 parts of sulphur.

Exp. 2. 1,175 parts of sulphuret of carbon produced 0,66 of carbonic acid gas, = 0,1782 parts (which is equivalent to 15,167 per cent.) of carbon; and 0,271 parts of sulphureous acid gas = 0,138 of sulphur. In the superior extremity of the glass tube which contained the oxyd of iron, there was found, in that part of the tube which projected out of the stove, a small portion of sublimed sulphur weighing 0,029. The sulphuret of iron found in the tube afforded, by the mode of estimation above related, 6,06 parts of ignited sulphate of barytes = 0,829 of sulphur; therefore the sum total of sulphur is 0,829 + 0,29 + 0,138 = 0,996 parts of sulphur, which added to 0,1782 parts of carbon, makes 1,1742 parts, or 0,0008 less than the weight of the sulphuret of carbon employed in the analysis.

B. *Comparison of the above Proportions of Sulphur and Carbon, in the Sulphuret of Carbon, with those which might be inferred from the laws of determinate Proportions.*

The law respecting the combination of combustible bodies between themselves, is that, *when two such bodies unite, the proportion in which this combination takes place is such, that if they*

be oxydated to a certain degree, they will either absorb an equal quantity of oxygen, or the one will absorb two, three, four, &c. times as much oxygen as the other. As to the metallic sulphurets in general, those which are called sulphurets *at the minimum*, are so composed that the sulphur requires twice as much oxygen in order to become sulphureous acid, or three times as much in order to become sulphuric acid, as the metallic body requires to become an oxyd or saline basis; and it is on that account, that the sulphurets in minimum produce, by their oxydation, neutral sulphites, or sulphates. Now, by comparing the quantities of oxygen required to convert the sulphur into sulphureous acid, and the carbon into carbonic acid, it appears that the first quantity is twice as great as the latter. The small difference is to be ascribed to the impossibility of obtaining perfectly accurate results in analytical processes so complicated as this; and therefore the sulphuret of carbon may be considered as constituted according to the same law as the other sulphurets; so that the sulphur which it contains requires twice as much oxygen to become sulphureous acid, as the carbon requires to become carbonic acid. Admitting, therefore, that the above views of the composition of those two acids are correct, the sulphuret of carbon would consist of 15,47 parts of carbon, to 84,53 parts of sulphur; and in that case the result of our analysis of the alcohol of sulphur would only differ by $\frac{1}{4}$ per cent. from the proportions obtained by that mode of estimation.

C. *Observations on the atomic doctrine of chemical Combination.*

Mr. DALTON has lately proposed a mode of viewing the subject of determinate proportions, which is distinguished by

its extreme simplicity. From that ingenious mode of expressing combinations, the sulphuret of carbon ought to consist of two atoms of sulphur to one of carbon, since, according to that system, both the sulphureous and carbonic acids are composed of two atoms of oxygen to one of radical. But it is probable, that the gaseous oxyd of carbon consists of two atoms of carbon to one of oxygen, because the quantity of carbon required to convert carbonic acid into carbonic oxyd, exactly doubles the volume of the acid. In this case, and if, according to Mr. DALTON's supposition, sulphureous acid consists of two atoms of oxygen to one of sulphur, the alcohol of sulphur would be, like all the other sulphurets at the minimum, composed of one atom of sulphur to one of carbon.

Sir H. DAVY, in his Elements of Chemical Philosophy, adopts, like Mr. DALTON, the idea that sulphureous acid gas consists of one portion of sulphur to two of oxygen; and agreeably to this, he supposes for instance that the sulphurets of copper, or of silver, are composed of one portion of sulphur to one of the metal. But on the other hand, he considers other sulphurets, such as those of antimony, of iron, and of lead, as consisting of two portions of sulphur to one of the metal, and the sulphuret of zinc would, according to his views, be composed of two portions of the metal to one of sulphur. Yet, notwithstanding this great diversity in the estimation of proportions, all these sulphurets are so constituted that, if the compound be fully oxydated, it remains in a state of perfect saturation. The cause of these apparent inconsistencies seems to be, that the supposed atom, or ultimate particle, or that which is to be regarded as the single portion of a body, requires to be fixed according to some determinate law, before Mr. DALTON's

ingenious method of expressing the proportions in which bodies combine, can be applied with certainty and precision.*

D. *Experiments on the Combination of the Sulphuret of Carbon, with the Alkalies, the Earths, and the metallic Oxyds.*

The following experiments will shew in an unequivocal manner, that the sulphuret of carbon is capable of combining with saline bases. These combinations constitute a new class of bodies, for which there is no name in our present chemical nomenclature. I shall propose to call them *carbosulphurets*, a name quite consonant with the principles of the received chemical nomenclature.

The unavoidable presence of water, in either caustic potash or soda, induced me to try, in the first instance, the action of ammoniacal gas on the sulphuret of carbon.

Carbosulphuret of Ammonia. Some ammoniacal gas, and some liquid sulphuret of carbon were successively introduced into a receiver filled with mercury, the sulphuret being enclosed in a small glass bulb having an open orifice. The first effect of this contact was a dilatation of the gas; the surface of the sulphuret soon covered itself with a pulverulent straw-coloured substance, as if a portion of sulphur had been precipitated. After a few hours, the gaseous mixture had sensibly

* It appears to me that the best way to form a system of definite proportions, and to make it harmonize with the general views of chemistry, would be to take oxygen as the base of the scale, because most chemical combinations turn upon the proportion of that ingredient; and as gaseous bodies unite in equal or multiple volumes, I would consider as *the atom* of any other gas, the proportional weight of an equal measure of that gas.

diminished in bulk, and as this diminution proceeded, the gas gradually deposited on the surface of the glass a saline yellowish substance, which did not exhibit, even with the aid of a microscope, the least appearance of crystallization. The mercury having at last filled the receiver, an additional portion of ammoniacal gas was introduced, and this was repeated till no further absorption took place. The whole of the sulphuret of carbon was then found converted into the yellowish uncrystallized matter just described. This substance had a strong smell of ammonia, and was so deliquescent that it could not be transferred from one vessel into another, without undergoing an obvious alteration. The solution of this substance in water is first red, but it very soon passes to a deep orange colour, shewing that it undergoes a partial decomposition; and if it be distilled in its solid, though humid state, it sublimes and deposits small shining crystals of hydrosulphuret of ammonia; whilst, on the contrary, if it be heated in the same vessel in which it is formed, that is, without any access of air or moisture, the carbosulphuret of ammonia sublimes unchanged, from one part of the vessel to the other, and no vestige of hydrosulphuret is perceived. It appears therefore, that the sulphuret of carbon can enter into combination with pure ammonia without depositing its carbon; but if moisture or air be admitted, an alkaline hydrosulphuret, or sulphuret is formed, and carbonic acid is generated.

Carbosulphuret of lime. If some pure quick lime be heated in a glass tube, by means of a lamp, and some sulphuret of carbon in vapour be made to pass through the heated earth, the latter becomes ignited at the moment the vapour comes into contact with it, and this ignition continues till the earth is

saturated. During this process, none of the sulphuret of carbon escapes, the whole of it being absorbed by the lime. The earth, at its surface, is found yellowish, owing to the formation of a little sulphuret of lime, but this appearance ceases on removing the surface, shewing that it arises from the contact of air. The earthy mass is tasteless, when first applied to the tongue, but a bitter taste, with a smell of sulphuretted hydrogen are soon perceived. It is no longer susceptible of being heated or dissolved by water. If digested with water in close vessels, a solution of hydrosulphuret of lime is obtained, and the undissolved portion is mostly found to be carbonate of lime.

Carbosulphurets of barytes and strontian may be produced in a similar manner, and are found to possess analogous properties.

There can be no doubt but that the two fixed alkalies, in a state of perfect dryness, would form, with the sulphuret of carbon, *carbosulphurets of potash and soda*, quite analogous to those just described; but as these alkalies cannot be obtained in an anhydral state, I could only examine their action on the sulphuret of carbon when in a state of solution. The general result of these trials was, that if a quantity of the sulphuret of carbon be long digested, at a very gentle heat, with a solution of caustic potash, the sulphuret of carbon is decomposed, and the mixture resolves itself into a hydrosulphuret, and carbosulphuret, and a carbonate of potash.

In the same manner, if sulphuret of carbon be long digested with liquid caustic ammonia, it dissolves very slowly, forming an orange coloured solution analogous to that in caustic potash just described.

I tried also to produce *carbosulphurets of metallic oxyds.* The process simply consisted in precipitating metallic solutions, by solutions of the sulphuret of carbon in caustic potash. The precipitates which were thus obtained, had characters sufficiently distinct to shew that they were chiefly metallic carbosulphurets; but as the presence of a hydrosulphuret, and of carbonate of potash, in the solutions used for these precipitations, necessarily interfered with the distinctness of the results, I shall not trouble the Society with a detail of these experiments.

E. *Experiments to determine the nature of a particular Substance, produced by the action of nitro-muriatic Acid, on the Sulphuret of Carbon.*

We have already alluded, Dr. MARCET and myself, in the course of our paper, to a new and singular substance, which we produced by the long continued action of nitro-muriatic acid on the sulphuret of carbon. This compound was discovered in the following manner.

A portion of sulphuret of carbon was exposed to the action of a mixture of fuming nitric acid with concentrated muriatic acid. The acid instantly acquired a peculiar odour resembling that of muriated sulphur. Upon trying to promote the combination by applying heat, the sulphuret of carbon was expelled in the form of vapour, which obliged us to leave the mixture to its own spontaneous action, under the temperature of the atmosphere, which was then at least 70°.* The sulphuret of carbon soon passed to a reddish orange colour, which it communicated to the acid; and during this change,

* In August, 1812.

nitrous gas, with a strong smell of muriated sulphur, was slowly evolved. After the space of a week, the liquor began to cover itself with a white crystalline substance, losing at the same time its colour. On stirring the mixture, these crystals fell to the bottom, where they were immediately dissolved by the remaining sulphuret of carbon. This, however, in proportion as it saturated itself with the new formed substance, gradually lost its colour, acquired a greater consistence, and at last, at the end of three weeks, it was entirely converted into a solid white crystalline body, having the appearance of camphor.

This substance being separated from the acid, and washed with cold water, presented the following characters: it was colourless; its smell resembled both that of muriated sulphur and of oxyd of osmium; its taste was both acrid and acid; it was very volatile, melted at a gentle heat, and sublimed without residue. In fact, this body very closely resembles camphor in its external properties. It is insoluble in water, but readily dissolves in alcohol and ether, from which it is precipitated by water; it is also soluble in the oils, whether fixed or volatile, with which it forms transparent solutions. The spirituous solution of this substance has a peculiar, disagreeable, and very acid taste. It reddens litmus paper, and dissolves zinc with the disengagement of an extremely fetid gas. From this solution, a spirituous liquor is obtained by distillation, which possesses the same smell, and leaves a residue of muriate and sulphate of zinc. When a pure solution of the peculiar substance in alcohol is submitted to distillation, the products are, first, some sulphureous acid gas, and then alcohol strongly impregnated with muriatic ether; after which the air of the

vessel is found to contain portions of sulphureous and carbonic acid. The peculiar substance, in its dry state, does not alter litmus paper; but it reddens it strongly if the paper be moistened. When exposed to the action of boiling water, this body volatilizes through the water, but the remaining liquor contains some muriatic and a little sulphuric acid. Water, therefore, has the power of decomposing, though very slowly, this substance. Indeed if it be left a long time in contact with a small proportion of water, the liquor becomes strongly acid. Liquid caustic potash assisted by heat, dissolves the peculiar substance without any disengagement of gas. This solution is colourless; when neutralized by sulphuric acid no precipitation takes place from it, which shews that the alkali decomposes the peculiar body. Sulphuric acid, however, produces a slight effervescence, and the liquor exhales a smell of sulphureous acid. On adding to it a solution of sulphate of silver, muriated silver is precipitated.

These results tend to shew, that the substance in question contains sulphur and carbon in combination with oxygen, that is in the state of sulphureous and carbonic acid; for otherwise, some carbon would have been precipitated, or some sulphuret or hydrosulphuret of potash would have been formed.

And again, a portion of the problematic substance being sublimed through ignited lime in a glass tube, it was absorbed by the lime without any vestige of sulphuret of lime being formed, or any carbon deposited. And a similar experiment being tried with ignited metallic iron, instead of lime, some muriate of iron and a sulphuret of oxyd of iron* were formed, and carbonic acid gas was disengaged. This last

* *Sulphure d'oxidule de fer.*

experiment in particular shewed clearly that the substance in question contains carbonic acid.

The existence of three acids united in this compound, being thus ascertained, their respective proportions remained to be determined. It was by an experiment similar to the last described, that this object was obtained. The outline of this analysis, a minute detail of which would unnecessarily prolong this communication, was as follows:

A portion of the compound, after being weighed with great accuracy, was sublimed through a tube containing some very fine spiral iron wire in a state of ignition, the weight of which was three times that of the compound submitted to analysis. A decomposition took place, the products of which were, as in the experiment above related, muriate of iron, sulphuret of oxyd of iron, and a mixture of the carbonic oxyd and acid gases. These being carefully examined by appropriate reagents,* so as to form an estimate of the quantities of the muriatic, sulphureous, and carbonic acids contained in the peculiar compound, the following ultimate result, respecting the nature and proportions of its constituent parts, was obtained, viz.

Muriatic acid - -	48,74
Sulphureous acid -	29,63
Carbonic acid (and loss)	21,63
	100,00

* The proportion of muriatic acid was estimated by dissolving the muriate of iron in water, and precipitating the acid by nitrate of silver; that of the sulphureous acid was ascertained by treating the sulphuret of iron with nitro-muriatic acid, and precipitating the sulphuric acid formed by muriate of barytes: from the known relation between the sulphureous and sulphuric acid, the quantity of the former was easily deduced. And, lastly, the proportion of carbonic acid was gained by reducing the gaseous mixture to the state of carbonic acid, and absorbing this by caustic potash.

From the mode of computation adopted by Mr. DALTON, this triple acid would be composed of two atoms of muriatic acid, to one of the sulphureous, and one of the carbonic acid. It is a remarkable circumstance that the proportions of sulphur and carbon which prevail in the sulphuret of carbon, no longer obtain in this compound; one of the atoms of sulphur being expelled during the formation of the triple acid, and converted into sulphuric acid, which is found in the nitro-muriatic liquor.

The reason why the combination in question does not take place on burning the sulphuret of carbon in oxymuriatic gas, is that this gas does not yield a sufficient quantity of oxygen to acidify the sulphur and carbon, which are to unite with the muriatic acid, so that either nitric acid must be added, or a portion of muriatic acid must be detached from the oxymuriatic gas, by the agency of water, or of some other oxydated body.

It will be necessary to give a name to this triple acid. The combination of the fluoric with the boracic acid has been called *fluo-boric acid;* but the word *fluo* does not express the degree of acidification of the radical, which would be of great advantage in systematic language. In the Essay on the Latin Chemical Nomenclature, which I published some time ago, I proposed the name of *acidum boracico-fluoricum.* Upon the same principle, the combination, of the muriatic and carbonic acid, discovered by Mr. J. DAVY, would be named *acidum carbonico-muriaticum;* the combination of the sulphureous and muriatic acid (should such a compound be discovered), would be called *acidum sulphuroso-muriaticum;* and our triple acid would be *acidum muriaticum sulphuroso-carbonicum,* a name inconvenient by its length, but perfectly consonant to the principles of Chemical Nomenclature.

XXIV. *On the Means of procuring a steady Light in Coal Mines without the danger of Explosion.* By William Reid Clanny, *M. D. of Sunderland. Communicated by* William Allen, *Esq. F. R. S.*

Read May 20, 1813.

THE many dreadful explosions of fire-damp, or inflammable air, which have occurred in the extensive and well regulated coal mines of this district, in the course of the nine years during which I have resided in the county of Durham, have often excited my most serious attention; and latterly these explosions have caused the death of so many industrious people, that no individual, possessed of common humanity, can look on the subject with indifference.

Though the improved methods of ventilation have been attended by many solid advantages to the proprietors of coal-mines, it is nevertheless worthy of remark, that the increased frequency of explosions clearly demonstrate, that ventilation, in this case, has been no preventive.

Ventilation undoubtedly supplies atmospheric air; but it cannot obviate those inundations of inflammable air, (if I may be permitted the expression,) which, rushing from the old workings and caverns of the coal mine, overwhelm every thing before them. It is evident that ventilation, even in its improved state, has afforded no relief whatever; and here the apparatus, which, *in the first instance*, I have the

honour to lay before the Royal Society, will be found to afford a good light, unaccompanied by danger.

It very frequently happens that accumulations of carburetted hydrogen gas, mixed with atmospheric air, take place in the wastes, or old workings of the coal mines, and though much precaution is used for keeping this inflammable air confined to its proper places by means of partitions and folding-doors, nevertheless when, by carelessness or accident, this air comes into contact with any ignited substance, an explosion generally takes place.

These explosions happen when the pit-men are occupied in hewing out the coal at the *workings*, should they chance to open a cavern of unmixed carburetted hydrogen gas. This gas for the most part being pent up in a condensed state, rushes forth from a chasm, and forming what is locally denominated a *blower*, it suddenly mixes with the atmospheric air of the mine, and surrounding the lights of the pit-men, an explosion follows, commensurate with the quantity of hydrogen gas, which is frequently very considerable.

It will be unnecessary to detail the phenomena of an explosion of inflammable air, as they are already sufficiently known; but I hope it will not be unacceptable to the Society to record a few of the more considerable explosions, which have occurred in the course of the last seven years, in this district alone, independently of those which have taken place in other parts of the kingdom within the same time. In the summer of 1805, an explosion happened at Hebburn colliery, by which thirty-two pit-men were killed, who left wives and children in a destitute state, to the number of one hundred and five. About the same time, a colliery at Oxclose blew up, by

which, I understand, thirty-eight men perished, leaving eighteen widows, and seventy children unprovided for.

Soon after this melancholy catastrophe, ten men perished at Killingworth by an explosion. And about the same time, seven men were instantly killed, and several severely wounded by an explosion at Fenton Park colliery. On the 25th of May, 1812, the colliery at Felling exploded, by which ninety-two persons were instantly destroyed, leaving forty-one widows, and one hundred and thirty-three children to the protection of the public.

And, upon the 10th of October last, the Harrington Mill pit exploded, by which twenty-three people were killed, and many others severely wounded and scorched.

Thus, in the short space of seven years, upwards of two hundred pit-men were deprived, most suddenly, of their mortal existence, besides a great many wounded; and upwards of three hundred women and children were left in a state of the greatest poverty and distress.

The great danger of these explosions, even when every precaution has been taken, is manifest by their frequency, and indeed it may be expected, that an explosion will take place by means of a lighted candle the instant that the hydrogen gas amounts to one-twelfth part of the atmospheric air present, and that a similar effect will follow at all proportions from one-sixth to one-twelfth.

When ventilation, by the methods in general use, is found insufficient to carry off the fire-damp, as it arises in coal mines, large pumps are employed at the top of the shaft for that purpose, which are worked by steam engines. So frequent and instantaneous are the changes in the proportions of inflam-

mable air, from accidental circumstances, that it would be impossible at all times to ascertain, by a chemical process, at all parts of the mine, when danger is impending, for frequently the greatest differences of proportions exist at the same time, in different parts of such extensive works as coal mines. In fact, the miners know, from the appearance of the light of their candles, when the proportion of hydrogen gas is such, as to threaten an explosion; hence they carefully watch each other's candles, that they may desist in time, and escape instant destruction.

The excavations of coal mines are much greater than they are generally supposed to be: in some collieries they are continued for many miles, forming numerous windings and turnings, along which the pit-men have frequently to walk for forty or fifty minutes before they arrive at the *workings*, during which time, as well as when at work, they have no direct communication with the surface of the earth, but are entirely at the mercy of their greatest enemy, the inflammable air. This circumstance first impressed me with the idea that the light, by which the pit-men were to work, might be insulated. I was well aware that no preparation of phosphorus could supply a sufficient light for the purpose; an observation equally applicable to the miserable scintillations of *steel mills* (as they are termed), which have often exploded the inflammable air of the coal mines.

I find it needful here to remark, that, as far as applies to myself, the idea of insulating the light, and also the plan which I have adopted for carrying this idea into effect, by the construction of the apparatus or lamp, are perfectly original. This lamp may be managed with the greatest ease by any

boy of common understanding. It is so strong, that should large pieces of coal fall upon it, they cannot in the least injure it. Nor is there any chance of its being upset by any accident, as it may be worked at the very bottom of the mine; and it is likewise conveniently portable.

The combustion of the candle, within the lamp, is supported by the ordinary atmospheric air of the coal mine, which is supplied by a pair of common bellows through a stratum of water below the candle; at the same time a portion of the air already in the lamp, is driven through another stratum of water above the candle, and thus the air supplied may explode within the body of the lamp, without communicating the effect to the air in the mine, however highly it might be charged with carburetted hydrogen gas.

The moment the air enters into the lamp, it comes in contact with the candle, and consequently, upon all occasions, a *small portion only* of the air can be exploded, instead of the whole contents of the lamp; by these means several obvious advantages are secured. The air passing in a brisk current from below upward, close by the candle, carries the snuff with it, so that the light is always clear and steady.

I may also remark that *wherever* a person can exist from a sufficiency of atmospheric air, this lamp will afford a safe and abundant light, from one candle only, for the space of five hours at least. This lamp will, in all probability, be found very useful in the powder magazines of ships of war, and of forts, as also in those places where gunpowder is manufactured; but this observation is merely thrown out for the opinion of those who are more conversant upon such subjects.

Fig. 1.



Fig. 3.



Fig. 2.



Description of the Lamp.

(See Plate XV.)

Fig. 1. describes the lamp as it appears when ready for use.

Fig. 2. A section of the lamp.

A. The body of the lamp constructed of copper or japanned iron, terminating in

B. A conical tube, which carries off the air (deprived of part of its oxygen by combustion) through the water in the cistern C.

D. Is a cistern containing water, in order to prevent the lamp from being over-heated.

E. The window of the lamp made of very thick glass.

F. The candle supported upon a tin stand.

G. A tube furnished with a cock, in order to bring the water to a level within the lamp.

H. A cistern containing water, which may be drawn off by the cock I.

K. A tube from the bellows which delivers air for the supply of the lamp, through the water in the cistern H.

Fig. 3. L. Spare bellows with an elastic tube M which may be adapted to a tube conveying pure atmospherical air, or to a gasometer. Forty gallons of atmospherical air will be sufficient to keep the lamp burning for one hour.

Fig. 1. N and O, two cocks to draw off the water from the cisterns C and D.

XXV. *On the Light of the Cassegrainian Telescope, compared with that of the Gregorian.* By *Captain* Henry Kater, *Brigade-Major. Communicated by the Right Hon. Sir* Joseph Banks, *Bart. K.B. P.R.S.*

Read May 27, 1813.

The Cassegrainian telescope from its first invention to the present time, has generally been considered to be merely the Gregorian disguised, and to possess no other advantages over it than the capability of being made shorter with the same magnifying power. This opinion, joined to the inconvenience of its inverting the object, has caused it to be thrown aside, perhaps too hastily, and without a sufficient examination of its properties.

As the experiments which I am about to detail may possibly lead to important conclusions, I shall perhaps be pardoned if I relate the circumstances which induced me to engage in them.

A self-taught artist of the name of Crickmore, who resides at Ipswich, had by exclusive attention to the subject, brought the Gregorian telescope to a degree of perfection surpassing any thing of the kind I have ever yet met with. Some months since, in the course of his experiments, he first completed a Cassegrainian telescope of one foot in length, and on viewing Jupiter with it, with a power of about 100, I was instantly struck with the brightness of the image, far exceeding what might have been expected from the aperture; but I supposed

this to have been a deception arising from the specula being more exquisitely figured than usual, which, producing greater distinctness, occasioned the idea of superior light. A short time after this, the same artist received an order for another telescope, which, from the success that had attended his recent efforts, he recommended to be of the Cassegrainian form. The aperture was five inches, the length thirty inches, and with a power of near 400 the image was so perfectly distinct and luminous, that I could no longer hesitate to conclude that, from some unknown cause, the Cassegrainian telescope actually possessed far more light than the Gregorian, and I waited most anxiously for an opportunity of verifying this, and determining the difference by experiment.

Such an opportunity soon presented itself, and under circumstances peculiarly favourable, as another excellent telescope of the Cassegrainian form was made, and I was fortunate enough to procure a Gregorian made by Mr. CRICKMORE some time before. The mirrors of both these telescopes were cast at the same time, and from the same pattern, so that no difference of light could arise from any difference in the composition of the metal. The magnifying power of both instruments was ascertained by experiment to be very nearly equal; but the excess was rather on the side of the Cassegrainian.

The telescopes being placed side by side, were directed to a printed card, at the distance of fifty yards; and on viewing it, the far superior brightness of the image in the Cassegrainian was strikingly apparent. Having prepared a circular piece of paste-board to close the end of the Cassegrainian telescope, I drew a number of concentric circles on it, at the distance of the twentieth of an inch from each other. The paste-board

was then placed in the end of the tube, and an aperture was made, which was enlarged by cutting out one circle after another till the card appeared equally bright through both telescopes, and of this, the eye judges most accurately.

The following measures were then taken.

Cassegrainian Telescope.	Inch.	Gregorian Telescope.	Inch.
Diameter of the circular opening in the paste-board	2, 70	Diameter of the large mirror	3, 90
Diameter of the back of the small mirror	1, 09	Diameter of the back of the small mirror	1, 00
Length of the arm	0,805	Length of the arm	1, 45
Thickness	0,200	Thickness	0, 20

From the above measures the following calculations were made.

Cassegrainian Telescope.	Inch.	Gregorian Telescope.	Inch.
Area of the circular opening in the paste-board	5,726	Area of the large mirror	11,946
Area of the back of the small mirror to be deducted	0,933	Area of the back of the small mirror to be deducted	0,785
Area of the arm to be deducted	0,161	Area of the arm to be deducted	0,290
	1,094		1,075
Area of the portion of the mirror exposed to the light	4,632	Area of the portion of the mirror exposed to the light	10,871

From this experiment it appears, that the light in both telescopes was equal when the area of the aperture of the Cassegrainian, was to that of the Gregorian, as 4,632 to 10,871. Now the increase of light being (under similar circumstances) directly as the area of the aperture, it follows that if the aperture of the Cassegrainian be made equal to that of the Gregorian, the light in favour of the former will be as 10,871 to 4,632, or in the surprising proportion of 7 to 3 nearly.

A difference of such magnitude could not be admitted but with extreme caution, particularly as the Gregorian telescope had been made some time, and its mirrors might therefore be supposed not to possess so high a polish, as those of the Cassegrainian which had been recently finished; but I was soon enabled to pursue the subject, as a Gregorian telescope was made by Mr. CRICKMORE fully equal, if not superior, to any he had before constructed; the mirrors were of an exquisite polish. The Cassegrainian, used in this experiment, was the one I formerly mentioned, the aperture of which was five inches, and the length thirty inches. It had not been carefully preserved, and the large mirror had lost somewhat of its original polish. All circumstances being thus in favour of the Gregorian, a paste-board circle was prepared, and the experiment conducted as before. When the images of the card were equally bright, the following measures were taken.

Cassegrainian Telescope.	Inch.	Gregorian Telescope.	Inch.
Diameter of the circular opening in the paste-board	3, 50	Diameter of the large mirror	3, 95
Diameter of the back of the small mirror	1,375	Diameter of the back of the small mirror	0, 35
Length of the arm	1,063	Length of the arm	1, 50
Thickness of the arm	0, 20	Thickness of the arm	0,175
		Length of a bar containing the adjustment	0, 70
		Its width	0, 15
		Diameter of three semicircles used as rests for the great mirror	0,375

From the above measures the following calculations were made.

Cassegrainian Telescope.	Inch.	Gregorian Telescope.	Inch.
Area of the circular opening in the paste-board	9,621	Area of the large mirror	12,254
Area of the back of the small mirror to be deducted	1,485	Area of the back of the small mirror to be deducted	0,709
Area of the arm to be deducted	0,213	Area of the arm to be deducted	0,263
	1,698	Area of the bar containing the adjustment to be deducted	0,105
		Area of the three semicircles to be deducted	0,166
			1,243
Area of the portion of the mirror exposed to the light	7,923	Area of the portion of the mirror exposed to the light	11,011

The magnifying power having been determined (by experiment) to be 188 in the Cassegrainian, and 182 in the Gregorian, the expression for the relative quantity of light becomes $\frac{11,011}{182^2}$ to $\frac{7,923}{188^2}$ or as 332 to 224, being nearly as 3 to 2.

In the first experiment, the advantages of polish were, *perhaps*, on the side of the Cassegrainian telescope; in the last, they were *much* in favour of the Gregorian; a mean therefore of both results may probably be considered as approaching the truth, and the light of a telescope of the Cassegrainian construction, may be taken, to that of a Gregorian of the same aperture and power, as about 60 to 33.

A fact so new, naturally leads the mind to hazard a conjecture as to the cause. In the Gregorian telescope a column of light from a point of the object, is received on the large mirror, and reflected in a cone of rays, the vertex of which is its focus, where an image is formed. Here all these rays meet in a single *point*, and crossing each other, fall on the small concave mirror whence they are again reflected, and form another image near the eye. Now, if light be supposed to consist of particles of matter, is it not possible that these particles, crossing in the same point, may interfere with each other? or, when thus forced within a certain distance of each other, may not a power of repulsion exist, which would occasion many of them to be dissipated? In the Cassegrainian telescope the rays reflected from the great mirror are received by the small convex mirror *before* they arrive at their focus, and are consequently reflected back without having crossed as in the Gregorian. The conclusion then seems to be, that wherever an image is formed, much light is lost, and this con-

clusion perhaps derives additional force from a circumstance noticed in most elementary works on optics, viz. that the satellites of Jupiter and his belts, may be distinctly seen with a Galilean telescope, whilst with an astronomical telescope of an equal aperture and power, they remain invisible.

Ipswich, 22d April, 1813.

XXVI. *Additional Observations on the Effects of Magnesia in preventing an increased Formation of Uric Acid; with Remarks on the Influence of Acids upon the Composition of the Urine.* By William Thomas Brande, *Esq. F. R. S. Prof. Chem. R. I. Communicated by the Society for improving Animal Chemistry.*

Read June 3, 1813.

In a paper which I had the honour of laying before this Society, about three years ago, and which is published in the Philosophical Transactions*, some cases are related, illustrating the effects of magnesia in preventing an increased formation of uric acid, and some experiments are detailed, instituted with a view to discover its mode of action.

Since that period many opportunities have occurred, both to Sir Everard Home and myself, of confirming its efficacy upon a more extended scale, and of ascertaining the efficient treatment of those cases in which magnesia is ineffectual, and in which it has even been found to aggravate the complaint.

To bring forward additional evidence in favour of the use of magnesia, and to distinguish the cases in which its use is indicated, from those where it is improper or hurtful, are the principal objects of the present communication, and will be considered in the two following sections.

* For 1813, p. 106.

SECTION I.

The following is the case of a gentleman who suffered from a calculous complaint, during which he was accidentally induced to employ magnesia, the effects of which he has thus described.

Case 1. About twenty-seven years ago, I felt a pain in one of my kidnies, particularly when in bed, which continued to increase during six months. I had likewise an occasional sympathetic pain in the testicles, and violent and excruciating pains in the left kidney now became frequent. These attacks were sometimes brought on by stooping to take up something; but at other times without any apparent cause. They lasted from twelve to twenty-four hours, and I obtained some relief from the application of warm flannels; but they always left me languid and relaxed.

On the fourth attack I consulted a physician, who imagined that my complaint had been induced by drinking cyder, in which I had formerly indulged. He ordered me weak Hollands and water for common drink, and prescribed the lixivium of tartar to be taken in broth. This medicine was persevered in for some time; but I found it gradually weaken my stomach, and impair my digestive powers.

About nine months after my first attack in the kidney, I walked from Hampstead to London after dinner, and on the following day, I clearly felt something pass from the kidney to the bladder, and suspected what it was. I took about a pint of Hollands and water, and on attempting shortly afterwards to void my urine, found that the passage was blocked up, but had scarcely time to consider of my situation before the

obstruction moved forwards to within an inch of the extremity of the urethra: it remained there till the following evening, when, by the help of a small pair of watchmaker's forceps, I succeeded in extracting a stone, which was the source of the mischief.

It was jagged and rough, and of a deep brick-red colour. I afterwards voided a considerable quantity of red crystalline sand.

My physician, who was apprehensive of a return of the disorder, desired me to purchase of CADELL, an anonymous pamphlet upon the Stone and Gravel, and to observe the rules there laid down. This treatise particularly recommended the use of the alkalies. I therefore took the lixivium, and two bottles of PERRY'S solvent; but the red deposit in my urine continued, my loins felt weak, and when in bed very painful.

Being in the profession of the law, and much employed, I was under the necessity of leading a very sedentary life, which so aggravated my tendency to bile and indigestion, that I seldom could get above two or three hours sleep.

With a view to alleviate these symptoms, and not with any idea of its being beneficial to the stone, I resorted to magnesia, which I continued with little intermission for eight months in the dose of a tea-spoonful or two, every evening before I went to bed. The long vacation coming on, I gradually took more exercise, and used the cold bath. The tone of my stomach, at the end of the period I have mentioned, was so far restored as to induce me to set medicine of all kinds aside, except when any food or drink disagrees, when I occasionally resort to the magnesia. Under such treatment, the weakness and pain in my kidney left me, and the red sand entirely disappeared. I

have since enjoyed a very good state of health, and am now in my fifty-seventh year.

If I occasionally make a little free with the good things of this world, my stomach reminds me of the improper use of the lixivium, especially when I am prevented taking my usual exercise.

The above case is important, not only as furnishing a striking and unprejudiced instance of the effect of magnesia, in counteracting the tendency to form uric calculi and gravel; but likewise, as demonstrating its efficacy where the alkalies had failed, and where the digestive organs had been injured in consequence of the use of such remedies: the time which has elapsed since the cure of this and other cases, without a relapse, is also strongly in favour of this mode of treatment.

Case 2. A gentleman twenty years of age who had suffered from heartburn, and other dyspeptic symptoms, was seized, on the 1st of June, 1811, with a violent pain in the loins, and more especially in the right kidney, and during the night he passed a large quantity of red sand with his urine. On the 2d, with a view to relieve the pain, which had increased considerably, he took fifty drops of laudanum, and drank freely of barley water. The night was passed more quietly, but on the morning of the 3d, he was seized with a violent pain in the kidney, and with the usual symptoms of the passage of a calculus along the ureter. These continued with more or less violence till the evening of the 4th, when he became perfectly easy, and remained so till the morning of the 6th, when, with

considerable pain and difficulty, he voided a calculus composed of uric acid, weighing nine grains. For several successive days his urine deposited a large quantity of red sand, and three very small round calculi were voided.

He was now directed to abstain from all kinds of fermented liquors and sour food, and to take a pint of treble soda water, (containing three drachms of sub-carbonate of soda,) daily. Under this treatment he continued to recover, and remained perfectly free from complaint until the end of August, when a copious deposit of red sand appeared in his urine: he had little pain in the affected kidney, but complained of almost constant nausea, or want of appetite. The soda water was increased to a pint and a half, and afterwards to two pints daily, and in the intervals he drank very freely of barley water.

Having persevered in this way for ten days without receiving any benefit, he was induced to make a trial of magnesia, of which he took one tea-spoonful night and morning in cold chamomile tea. In about a week, the state of his stomach was much improved, and the deposit in the urine proportionally diminished, and in three weeks every symptom of disease had disappeared.

In February, 1819, having persevered in the use of magnesia with little intermission, I was informed that the sand had returned, that increasing the quantity of magnesia had produced no good effect, and that alkalies materially aggravated his complaint, by disagreeing with the stomach and greatly increasing the urinary deposit.

On examining the sand, I found that instead of consisting as formerly of uric acid, it was composed of a mixture of the

ammoniaco-magnesian phosphate with phosphate of lime; he was directed to abstain from magnesia and alkalies, and to adopt a plan of treatment which it is the object of the second section of this paper more particularly to explain.

The foregoing is a well marked case of uric gravel with a strong tendency to form calculi, materially relieved by the use of alkaline remedies: it illustrates their usual effects when carelessly persevered in, and shews the advantage with which magnesia may in such instances be employed: it also exhibits the effect of magnesia and the alkalies, in producing the deposit of *white sand* (or phosphates) in the urine, when the *red sand* (or uric acid) has been removed.

The cases which follow are selected, from among others, to explain the best mode of preventing the formation of white sand, and to shew the most effectual treatment where it is a natural deposit in the urine, or where it has been induced by the incautious exhibition of alkaline medicines.

SECTION II.

The white sand so frequently voided by persons labouring under calculous complaints, was first analyzed by Dr. WOLLASTON,* who found it composed of ammoniaco-magnesian phosphate, either alone or mixed with variable proportions of phosphate of lime. The use of acid medicines in these cases was also first suggested by the same able chemist, but although his valuable observations have been before the public for nearly fifteen years, I am not aware that any

* Phil. Trans. 1797.

experiments have been made to ascertain what acids are best calculated to produce the desired effect, or to illustrate their mode of action.

Since my former communication, I have lost no opportunity of attending to this important subject, and hope that the conclusions, suggested by the following cases, will be deemed satisfactory, and that their application in practice may lead to useful results.

Case 1. A gentleman, fifty years of age, who about ten years before had undergone the operation for the stone,* was attacked on the 14th of January, 1810, with violent pain in the right kidney and ureter, which lasted two days; on the 17th, these symptoms subsided, and were followed by those of stone in the bladder, which continued for some days, and although he had taken abundance of barley water and similar diluents, the stone shewed no disposition to pass. On account of his former sufferings, this circumstance rendered him extremely uneasy, and on the evening of the 21st, he suffered several severe paroxysms of pain on attempting to make water. Under these circumstances, he was desired to take a purge, composed of two ounces of infusion of senna, two drachms of tincture of senna, and twenty grains of powdered jalap.† In three hours this began to take powerful effect, and during the

* The stone extracted consisted of a nucleus of uric acid about the size of a pea, incrusted with a mixture of the phosphates. It was broken during the operation, but appeared to have been of the size of a pigeon's egg.

† I recommended this treatment in consequence of having heard Sir Everard Home state a case, in his Surgical Lectures, of a gentleman who suffered a bougie to pass so far into the urethra, that it could not be removed by any instrument. During the operation of a purge it was expelled with considerable force.

violence of the operation, he was so fortunate as to void the calculus with his urine; it weighed eight grains. On the 28th he again suffered pain in the region of the kidneys, and voided much sand, composed of uric acid, with ammoniaco-magnesian phosphate. He now took three half pints of soda water daily, which materially increased the proportion of the triple phosphate, while that of uric acid was considerably diminished. Ten drops of muriatic acid were then taken three times a day in water. The red sand now began to re-appear, and on the 4th of February, he voided a very small uric calculus. The urine made after dinner contained more or less mucus streaked with blood, a symptom which was much aggravated by a slight excess in wine. On the 6th, he left London, and employed no medicine until the 12th, when he returned in consequence of having voided a large quantity of the white sand.

Having observed the efficacy of carbonic acid in preventing the deposition of the phosphates, and having found it less liable than any other acid to induce a return of the uric gravel and calculi, I now directed him to take half a pint of water highly impregnated with fixed air, four or five times a day, and to drink cyder instead of wine. On the 18th of February, his urine was less turbid than it had been for some months before, and on the 20th of March, having continued the use of carbonic acid, he had no remaining symptoms.*

* I have several times examined the urine, with a view to ascertain whether any of the acids which were exhibited, could be detected in that secretion; but the results of such experiments are so much interfered with by the very compound nature of the urine, that I have not hitherto been able to draw any satisfactory conclusions respecting them.

In August his urine became again turbid, but by the use of vinegar and lemon juice at his meals, which acids, he now finds, have no tendency to induce a return of the red gravel, he succeeds in preventing this symptom.

Case 2. On the 11th of October, 1812, the operation for stone in the bladder was performed upon a boy, eleven years of age, and four calculi were extracted, of which the largest was of the size of a small horse bean: they were each composed of a nucleus or centre of uric acid, upon which the ammoniaco-magnesian phosphate was deposited.

After the operation, the urine deposited a large quantity of white sediment, and some small pieces of red gravel were occasionally voided. He was now directed to take eight grains of citric acid dissolved in barley water, three times daily; under this treatment the sediment in the urine was considerably diminished, but did not wholly disappear. The dose of the acid was gradually increased to twenty grains, by which means the sediment was only occasionally deposited, and consisted of little else than mucus. It was observed, that whenever the citric acid was omitted, even for twenty-four hours, the sediment was greatly increased, and this was constantly attended with frequent desire to make water, and other symptoms of irritation in the bladder. On resuming the use of the citric acid, the sediment always disappeared, and the irritation of the bladder subsided, and this happened so frequently, that no doubt could be entertained of the influence of the medicine on the composition of the urine.

This plan of treatment was continued for three months; at the end of that period, it was found that the urine had not the

same disposition to deposit the phosphates as formerly; even when the medicine was omitted, the sediment was small in quantity, and not constant in its appearance. He was now directed to omit the use of the citric acid, and occasionally to eat oranges and other acid fruits. He continued this plan until the beginning of April, 1813; his urine was then quite clear, and he had no symptoms of disease.

Case 3. In the month of October, 1811, a gentleman, thirty-four years of age, informed me, that he had observed a white deposit in his urine, during the whole of the preceding summer. He had taken considerable quantities of soda water, which he thought increased the sediment, and alkalies in any other form produced a very obvious aggravation of the complaint.

His urine was at all times clear when voided; but after a few hours, a white powder was observed to separate from it, and a film of crystalline matter formed upon the surface. The former consisted of phosphate of lime and mucus, the latter of the ammoniaco-magnesian phosphate.

He was directed to take one drachm of muriatic acid properly diluted, at divided doses, during the day; and it was proposed that he should pursue this plan for a week; but it was discontinued on the third day on account of its acting upon the bowels, and producing a frequent desire to make water.*

On the 10th of October, he was advised to take two large glasses of lemonade daily, and to substitute claret for port wine, a pint of which he was in the habit of drinking daily.

* In this and other instances the sulphuric and nitric acids were occasionally substituted for the muriatic; but they were found equally inadmissible.

Under this treatment the symptoms produced by the muriatic acid subsided; but the appearance of the urine was not at first improved.

On the 20th, the film of triple phosphate formerly constantly observed in the urine began to decrease, but the white sand remained as abundant as before; he was therefore directed to take twenty grains of citric acid twice a day, and to continue the use of acid drink, as formerly.

The additional acid at first disagreed with the bowels; but this effect soon ceased, and the sediment was only observed in the urine voided in the morning; he therefore took another dose of the acid every night. This plan was pursued with little intermission until the beginning of December: the deposition of the phosphates gradually ceased, and he remained in perfect health until the middle of May, 1812, when after violent exercise and taking more wine than usual, the white sand again made its appearance in great abundance; his stomach became extremely irritable, and the acids, which he had before employed with success, brought on considerable irritation in the bladder. The addition of ten drops of laudanum to each dose of the citric acid prevented this effect, and he was thus enabled to continue the acid, which in a fortnight relieved his complaint.

This gentleman informed me, that whenever he omitted the use of an acid diet, or took much wine, especially port, his urine deposited the white sand and mucus, for two or three successive days.

Case 4. A gentleman, eighty years of age, who had twice submitted to the operation for the stone within five years,

voided with his urine considerable quantities of white sand and mucus.

From the age of this patient, and the account of his case, there appeared little doubt that the calculi had been formed in consequence of a diseased prostate gland, in the manner described by Sir EVERARD HOME,* and on examining them, they were found to contain no uric nucleus, nor indeed had there been any symptoms of disease in the kidneys, at any previous period.

This gentleman had been in the habit of taking soda water, from which he was now desired to abstain, with a view of putting him upon the acid plan of treatment. He was ordered to take eight drops of muriatic acid three times a day in two table-spoonsfull of water; but the third dose produced so much irritation in the bladder, and consequent increase of his symptoms, that it became necessary to adopt another treatment.

Lemon juice, or a solution of the pure citric acid, when given in quantity sufficient to produce any change in the appearance of the urine, had the same effect as the muriatic acid.

As water impregnated with carbonic acid could not be procured, he was directed to dissolve, in separate portions of water, twenty grains of citric acid, and thirty grains of the crystallised carbonate of potash, and to take the mixed solutions, during the effervescence. This quantity was at first only taken night and morning, but as it agreed perfectly well, it was afterwards repeated four and five times daily. Under these circumstances the appearance of the urine was soon improved, and both the mucus and the sand were considerably

* Practical Observations on the Treatment of Diseases of the Prostate Gland, p. 39.

diminished in quantity. In six weeks the urine, when voided, was transparent; but a considerable deposition of the phosphates took place, when it had remained for some hours at rest. In this state he left London, and has since informed me, that the sediment gradually diminished under the use of the carbonic acid, that his urine is never turbid, and that the irritation in the bladder has entirely subsided.

It did not appear necessary to detail the minutiæ of the above cases; they have been selected with a view to elucidate the treatment of the disease, as far as it depends upon chemical principles, and to furnish the data upon which the following conclusions are founded.

1. That where alkalies fail to relieve the increased secretion of uric acid, and to prevent its forming calculi in the kidneys, or where they disagree with the stomach, magnesia is generally effectual, and that it may be persevered in for a considerable time without inconvenience, where the tendency to form excess of uric acid remains.

2. When the alkalies, or magnesia, are improperly continued, after having relieved the symptoms connected with the formation of the red sand, or uric acid, the urine acquires a tendency to deposit the white sand, consisting of the ammoniaco-magnesian phosphate and phosphate of lime.

3. The mineral acids, (muriatic, sulphuric, and nitric) diminish, or entirely prevent the deposition of the phosphates; but are apt to induce a return of the red gravel.

4. That vegetable acids, especially the citric and tartaric, are less liable to produce the last mentioned effects, even when taken in large doses for a long time; and that carbonic acid is particularly useful in cases, where the irritable state of the bladder prevents the exhibition of other remedies.

XXVII. *Additions to an Account of the Anatomy of the Squalus Maximus, contained in a former Paper; with Observations on the Structure of the Branchial Artery.* By Sir Everard Home, *Bart. F. R. S.*

Read June 24, 1813.

My former account was taken from a Squalus Maximus caught at Hastings, in November, 1808, and the parts which I examined were brought to London by Mr. Clift, who went down, at my desire, to dissect them; but the weather being stormy and cold, the fish was brought no further than the beach, so that the examination was conducted under great disadvantages, and the parts brought away were in a mutilated state. The sketch of the fish made upon the spot by Mr. Clift is now found to be generally correct, except the omission of a small fin between the anus and tail, which had been buried by the weight of the fish in the sand.*

Two fishes of the same species have since been caught at Brighton, and one of them was brought to London in December, 1812, which I had an opportunity of examining, assisted by Mr. Clift. It is not my intention, on the present occasion,

* The omission of this small fin in the drawing is an error of considerable importance, as it deprived the fish of one of its characteristic marks, and has led naturalists, who have since had the opportunity of examining other specimens with more accuracy, to conclude that this fish was a distinct species from those which they described; I am therefore particularly desirous to correct the mistake.

to enter into minutiæ, but to render my former account more complete, and apply the dissection of this large fish to its proper use in comparative anatomy, which is, by means of it, to illustrate the functions of the organs of fishes of an ordinary size.

In addition to my former description of the fins, I have now annexed a drawing (Pl. XVI.) which shews the structure of the pectoral fin, a beautiful example of the mechanism of the fins of cartilaginous fishes in general.

The stomach was examined in its entire state, and the annexed drawing (Pl. XVII.) is an accurate representation of the appearance of its internal surface, which is exposed in one view, shewing that the pyloric portion is longer and narrower than I had before represented it.

The situation of the pancreas is correctly noticed in my former paper; the gland is oblong, thick and round where it is attached to the duodenum, and becoming thin, flat, and bifid towards its loose extremity.

The ducts of the liver are six in number, and inclosed in a broad flat band, which passes obliquely down before the stomach, till it is connected to the duodenum; each of the ducts opens, by a separate oblique orifice, into a common cavity of an oval form, from which there is a direct opening into the duodenum. This swell or enlargement might be considered as a substitute for the gall-bladder, which is wanting, were it not that a similar enlargement is also met with in fishes which have one. In the cod there is the same dilatation, and the hepatic ducts open into it in the same oblique manner; but there is also a gall-bladder, and the cystic duct, as well as the others, terminates in this dilatation.

The oblique openings of the hepatic ducts in the Squalus

Maximus, being so different from those of quadrupeds, explain the general principle on which the hepatic ducts in fishes, whose livers are loaded with oil, are formed. The substance of their liver is so exceedingly tender, that this contrivance is employed to prevent the bile from being forced back into the liver, which is not found necessary in the solid livers of land animals.

The heart was particularly examined, and the annexed drawing (Pl. XVIII.) shews its internal cavities, and the valves of the branchial artery; more particularly a muscular structure met with in the coats of that vessel, extending for some way after it leaves the ventricle.

The situation of the kidneys is mentioned in my former account; the ureters open into the cavity, common to the urine and semen, by two orifices, three quarters of an inch in diameter.

The structure of the body of the testicles had been destroyed; the epididymis consisted of innumerable convolutions of a tube three-eighths of an inch in diameter, at the upper part coiled up into two or three lobes or masses, which could not be unravelled, from which the vas deferens went off, making irregular convolutions down towards the anus. The lower part for three feet in length is straight, and the canal seven inches in diameter, having broad valvulæ conniventes; the widest are near the termination, and are two inches and a half broad. The contents of this portion, as I have remarked in my former paper, are different from those of the epididymis, and upper portion of the vas deferens, being a substance like starch broken down into rounded portions in a thinner fluid.

This circumstance leads to the idea, that the straight portion

of the vas deferens is analogous to the vesiculæ seminales of other animals, secreting its own fluid with which the semen is to be mixed before it enters the penis, and the valvulæ conniventes form a surface from which this secretion probably takes place.

It is deserving of observation, that the epididymis and vas deferens were loaded with semen, which there is reason to believe is not the case in quadrupeds, unless immediately previous to the act of copulation; but in fishes, where the connexion between the sexes is less complete, the semen appears to be prepared at a more early period.

The openings from the vasa deferentia are situated on each side, four inches higher up than those of the ureters, rounded in form, and kept closed by the pressure of two oviform ligamentous substances; these openings readily admit of dilatation, so that the hand can pass into them.

The penis has an infundibular form, and terminates by an oblique aperture about three inches in circumference.

The holders correspond with those of the dog-fish, which upon another occasion have been described to the Society; the spur bears a striking resemblance to that of the male ornithorhynchus paradoxus. There is a canal in each holder communicating with a corresponding cavity between the skin and muscles of the abdomen, eleven feet long and two wide. The inner surface of this cavity is smooth, almost polished, and of a beautiful white colour; it contained a white mucus, extremely viscid and tenacious.

Opportunities of examining the brain of a fish of such magnitude are of rare occurrence; I have therefore not only given drawings of the brain, (Pl. XIX, XX.) but also one

of the brain of a Squalus Acanthias, (Pl. XXI.) that the two may be compared together, with a view to shew the relative size of the parts, one belonging to a shark of thirty feet, the other to one of three feet long.

In the brains of what we consider the animals of greatest intelligence, there is a cerebrum, cerebellum, and medulla oblongata; beneath the cerebrum there are the tubercula quadrigemina. In fishes, the cerebrum is wanting, and there is no part at all analogous to it, unless we consider the enlargements, from which the olfactory nerves arise, to be of that description.

These enlargements are separated from the other parts of the brain by the optic nerves going off in a transverse line between them and the tubercula quadrigemina. In the present specimen, unfortunately, not only the enlargements from which the olfactory nerves go off, were destroyed; but also a portion of each of the anterior tubercula quadrigemina: the cerebellum was, however, entire, and is represented of the natural size.

The brain does not occupy more than one-third of the cranium. The medulla spinalis is large in proportion to the brain. From the termination of what corresponds to the calamus scriptorius in the human brain, a fissure extends on the upper part of the medulla spinalis, dividing it longitudinally into two portions; there is a similar fissure on the anterior surface; into both of these a thin fold of pia mater extends and adheres with firmness to the surfaces with which it comes in contact.

The dura mater is very dense, and adheres firmly to the inner surface of the cranium and theca vertebralis. The pia

mater nearly resembles that of the human brain; it becomes thicker where it covers the spinal marrow. The space between the dura and pia mater is occupied by a cellular membrane of a very fine texture.

As the different parts of the brain are described in the explanation of the drawing, I shall only remark in this place, that the circumstance most deserving of observation respecting it is, that the cerebellum has an increase of size in the Squalus Maximus in a much greater degree beyond that of the Squalus Acanthias than the tubercula quadrigemina. The protuberances from which the olfactory nerves arise were probably large in the same proportion with the tubercula; at least, in the brain of a shark, preserved in the Hunterian Collection, of a smaller size than that of the Squalus Maximus, but much larger than that of the Squalus Acanthias, that is the case.

The eye is small for the size of the fish; the ball has projections on the sclerotic coat, where the muscles are attached, which make it approach to a quadrangular form; but its internal cavity is circular. The circumference in the widest part is nine inches. The longest diameter three inches, the shortest one inch and three quarters. The sclerotic coat is cartilaginous, one quarter of an inch thick on the posterior part, becoming thinner towards the ciliary processes, where it is only one-sixteenth of an inch.

The cornea is thin, but made up distinctly of three layers, of which the middle one is by much the thinnest. The optic nerve is nearly of the size of the sixth pair, and, where it perforates the sclerotic coat, projects a little before it gives off the retina, which is extremely thin.

The choroid coat is covered with a tapetum lucidum of the

colour of an amalgam of silver broken into small portions. There are ciliary processes, not common to fishes in general; they are about one-third of an inch in extent, and slightly projecting; they are lined with a black pigment.

The vitreous humour is unattached to the choroid coat; it is inclosed in strong cells, and the crystalline lens, which is spherical and one inch in diameter, is imbedded in it for two-thirds of its substance. In the Tetrodon Mola, which, as well as the Squalus Maximus, in common language has been called the Sun-fish, the vitreous humour has a firm attachment to a groove in the choroid coat one-twelfth of an inch in breadth, extending from the entrance of the optic nerve to the termination of the retina, in the shortest line from the one to the other, and there are no ciliary processes; two such remarkable differences in eyes of nearly the same size, appeared to be deserving of being noticed.

The cartilage upon which the ball of the eye rests is attached to the bottom of the orbit, and is seven inches and a half long; its stem is a flattened cylinder, five-eighths of an inch in diameter; it terminates in a broad concave surface in a transverse direction adapted to the bottom of the ball of the eye, and is connected to the sclerotic coat by a ligament long enough to admit of motion.

There are four straight and two oblique muscles; the straight very large, much beyond what can be required merely to move so small an eye: the rectus internus and externus are strongest, they are five inches in circumference, while the superior and inferior are only three and a half.

In the structure of the ear, the only remarkable circumstance is the great capacity of the cavities in which the semi-

circular canals are contained, the canals themselves not being much larger than in the skate. They are shewn in the drawing. In the recent subject they were pellucid, containing a transparent fluid.

On the Structure of the Branchial Artery.

The muscular structure of the branchial artery of the dog-fish, and the direction in which that artery leaves the ventricle, are exactly the same as in the Squalus Maximus, only are seen upon so small a scale, that they do not arrest our attention; but when magnified, to the size which they acquire in this fish, they make a stronger impression on the mind, and force us irresistibly to enquire into their use.

This direction of the artery appears to be common to fishes in general; but the muscular structure is confined to particular tribes. I find it is common to all the sharks, and there is a similar structure in the sturgeon.

In the wolf fish, the Anarchichus Lupus, the muscular structure of the branchial artery is nearly the same; but the valves are placed close to the opening of the ventricle, and are only two in number.

In the turbot there is no muscular structure in this part of the artery; but the coats are extremely elastic, and admit of being very considerably dilated, particularly at its origin, where three valves are placed, and so contrived that the dilatation of the artery makes them shut more closely.

In the Lophius Piscatorius, there is no appearance of muscularity in the coats of the branchial artery, and no lateral valves as in other fishes; but there is a muscular tube half

an inch long, rising from the edge of the opening of the ventricle which projects into the artery.*

These different structures, so very unlike one another, and bearing no resemblance to the mechanism of the same parts in quadrupeds, make it probable, that the circulation through the gills is impeded by the external pressure of the water, in different degrees according to the depth of the fish from the surface: therefore in those fishes which frequent great depths, as the Squalus Maximus and all the shark tribe, there is a muscular structure in the coats of the branchial artery, which, when the fish is deep in the water, by its contraction diminishes the area of the vessel, and makes the valves perform their office; but when the fish is near the surface, this muscular structure, by its relaxation, renders the area of the artery so wide, that regurgitation of the blood takes place into the ventricle, and prevents the small vessels of the gills from being too much loaded.†

In fishes that swim deep, and do not come to the surface, as the wolf fish, the regurgitation does not take place into the ventricle; but the relaxation of this muscular portion of the artery allows it to dilate, and form a reservoir, and the valves remain closed, so as to prevent more blood leaving the ventricle. In fishes residing at moderate depths, as the turbot, elasticity is employed as a substitute for muscular power, there being less variation in the pressure made upon the gills; but in the Lophius Piscatorius, which probably never descends into

* This structure, which it is difficult to describe, will be better understood by a reference to the annexed drawing. (Pl. XXII.)

† That such regurgitation takes place when the muscle is relaxed, is ascertained by the ventricle being readily injected after death with common wax injection from the artery, the valves allowing it to pass.

water of much depth, the ventricle is so weak, that the supply of blood to the gills is regulated by the contraction and relaxation of a muscular valve.

As water, according to the degree of pressure upon it, is capable of containing a greater quantity of atmospherical air, than under ordinary circumstances, such a supersaturated state of the water might compensate, with respect to the respiration of fishes, for the difficulties, which occur at great depths, of forcing the blood through the vessels of the gills: I enquired what evidence could be produced, of the water at great depths containing a more than ordinary quantity of air; my philosophical friends, to whom I proposed this question, said, that it was a point that had not been considered. I therefore resolved to put it to the test of experiment, and as I knew there was a well at Mr. COUTTS's in the Strand, which more than twenty-five years ago had been sunk four hundred feet below the surface of the water in the Thames, I requested permission to make the experiment in that well. A cylindrical vessel, with a valve above and below, was let down to the depth of one hundred and eighty-six feet, which is now the depth of the well; it came up full of clear water, which Mr. W. BRANDE ascertained to contain no greater proportion of atmospherical air, than is met with in common river water. It therefore appears that the supply of air to fishes is nearly the same at whatever depth they are from the surface.

Besides the guards which have been mentioned, to prevent the circulation through the gills from being improperly carried on, in particular tribes of fishes, the ventricle itself is so formed, and is so situated with respect to the auricle, that the blood received is first impelled in a direction nearly at right

angles to that of the artery, and then another set of muscles is employed to force it into the artery.

This mechanism, common to most fishes, and so different from what is met with in land animals, where the blood is forced by the action of the left ventricle into the different parts of the body, appears to be intended to prevent too great a force being employed, at any one time, in impelling the blood through the gills.

Having traced in these different gradations the guards upon the circulation through the gills in fishes, I am led to extend my remarks to the hearts of a lower class of animals. In the Mollusca there is no reason for putting guards upon the action of the heart, as in fishes, because in them the blood first supplies the different parts of the body, and only in its return passes through the gills; but there is a regulation of another kind, by means of which the circulation is increased or diminished, according to the activity or torpor of the animal. In the teredines, where a boring engine, requiring a great muscular power to work it, is almost constantly employed, the heart consists of two auricles and two ventricles; the auricles strong, cylindrical, and having valves between them and the ventricles; the ventricles themselves very strong, so that in fact there is a heart, composed of two auricles and two ventricles, both acting at the same time, and the blood is hurried on by a double power to supply the muscles of the boring shells, and, in this part of its course, has a bright red colour.

In the Oyster, on the other hand, the heart consists of only one auricle and one ventricle, both very weak in their muscular power, when compared with those of the teredines, although belonging to an animal of larger dimensions; but the heart is

laterally connected to the great muscle that shuts the shell, so that whenever that muscle is actively employed, its lateral swell presses against the heart, and forces out the blood, and by this means gives activity to the circulation, at the only time in which an increased circulation is particularly required.

In the Muscle, the heart can hardly be said to be distinctly divided into auricle and ventricle; but to consist of an oval bag, through the middle of which the lower portion of the intestine passes. Two veins from the gills open into the heart, one on each side, these may be considered as the auricles.

The coats of the ventricle are very thin, so as to have little power of propelling the blood; but this weakness appears to be compensated by the effect of the contraction and relaxation of the intestine contained in their cavity. This circumstance renders it probable, that the circulation goes on with activity while the processes connected with digestion are employed; but when the intestines are empty, there being no supply of nourishment, the circulation is not only very languid, but may possibly be entirely stopped.

In the caterpillar, the blood cannot be said to circulate, but is carried from one end of the animal to the other, in a tube which may be either called heart or artery, by a species of peristaltic motion. This ebbing and flowing of the blood, if such a term can be admitted, is greatly increased by the pressure upon the different portions of this tube, produced by the action of muscles employed in the progressive motion of the animal, so that the supply of blood to the different parts of the body is proportioned to the demand, which arises out of the bodily exertions.

Fig. 1.

Two Inches to a Foot

Fig. 2.



Natural size

One Inch to a Foot.

Fig. 1.



Fig. 2.



2 Inches to a Foot.



J. Hawkins del.

F. Basire sc.

J. Rowlandson, del.



J. Basire sc.



J. Rowlandson del.

H. Meyer sc.



W. Clift del.

J. Basire sc.

EXPLANATION OF THE PLATES.

(See Plates XVI, XVII, XVIII, XIX, XX, XXI, XXII.)

Six of the following Plates are taken from drawings made by Mr. HOWSHIP, who very kindly undertook that task, during the indisposition of Mr. CLIFT, and his knowledge of anatomy, which he has cultivated with much ardour, induced me to solicit his assistance upon this occasion.

PLATE XVI.

Fig. 1. A view of the pectoral fin of the Squalus Maximus, from which the skin and cellular membrane are removed, to shew the arrangement of the cartilages of which it is composed.

Fig. 2. A portion of the elastic fibrous structure with which the fin is tipped, of its natural size.

PLATE XVII.

Fig. 1. An internal view of the stomach and duodenum of the Squalus Maximus.

a. The œsophagus.

b b. The cardiac portion of the stomach.

c c. The pyloric portion.

d d. The spleen.

e. A small cavity belonging to the stomach.

f f. The duodenum.

g. The band containing the hepatic ducts, six in number.

h. The dilatation in which the gall ducts terminate.

i. The opening of the pancreatic duct.

k. The spiral turns of the intestine.

Fig. 2. The fringe at the termination of the œsophagus of the natural size.

Plate XVIII.

The heart of the Squalus Maximus laid open.

a a. A portion of the internal surface of the auricle.

b b. One of the venæ cavæ laid open.

c. The valve between the auricle and ventricle.

d d. The cavity of the ventricle.

e e e. The three rows of valves and the three intermediate spaces, along which regurgitation takes place when the canal of the artery is dilated.

f f. The strong muscular covering of the artery.

Plate XIX.

A side view of the cerebellum, tubercula quadrigemina, and nerves of the brain of the Squalus Maximus.

3, 4, 5, 6, 7, 8. The different nerves going off from the brain corresponding with those in the brain of man.

a. Cerebellum.

b b. Two of the tubercula quadrigemina.

c. A third tuberculum mutilated.

Plate XX.

A view of the upper surface of the cerebellum and nerves of the Squalus Maximus.

The nerves going off marked, as in the last Plate, beyond which are two pair of nerves belonging to the spinal marrow.

a a. Cerebellum.

b b. Two of the tubercula quadrigemina.

c c. The part which corresponds with the fourth ventricle in the human brain, surrounded by an oval, continued, nervous band, from which the principal nerves go off.

d d d. Portions of the three semicircular canals of the ear in the cartilaginous cavity in which they are contained.

Plate XXI.

A view of the upper surface of the brain of the Squalus Acanthias, taken from a fish three feet long, to shew the difference of appearance and size between it and that of the Squalus Maximus.

The brain is entire, and the eyes are left in their situation, so that when this Plate is compared with that of the large brain, the parts that are wanting in it will be readily distinguished.

Plate XXII.

A view of the heart of the Lophius Piscatorius in a distended state, shewing the transparency of its coats, which are extremely thin.

a. The auricle.

b. The ventricle.

c. The branchial artery.

d. The projecting muscular tube serving as a valve.

XXVIII. *Some further Observations on a new detonating Substance. In a Letter from Sir* Humphry Davy, *LL.D. F.R.S. V.P.R.I. to the Right Hon. Sir* Joseph Banks, *Bart. K.B. P.R.S.*

Read July 1, 1813.

My dear Sir,

Berkeley-square, June 20, 1813.

I have already described, in a letter which you were so good as to communicate to the Royal Society, a few facts respecting a new detonating compound. I shall now do myself the honour of mentioning to you some other particulars on the subject.

I received, in April, a duplicate of the letter in which the discovery was announced, containing an Appendix, in which the method of preparing it was described. M. Ampere, my correspondent, states that the author obtained it by passing a mixture of azote and chlorine through aqueous solutions of sulphate, or muriate of ammonia. It is obvious, from this statement, that the substance discovered in France, is the same as that which occasioned my accident. The azote cannot be necessary; for the result is obtained by the exposure of pure chlorine to any common ammoniacal salt.

Since I recovered the use of my eyes, I have made many experiments on this compound; it is probable that most of them have been made before in France; but as no accounts of the investigations of M. Dulong on the substance have

appeared in any of the foreign journals which have reached this country, and as some difference of opinion and doubts exist respecting its composition, I conceive a few details on its properties and nature will not be entirely devoid of interest.

I have been able to determine its specific gravity, I hope, with tolerable precision, by comparing its weight at 61° FAHRENHEIT, with that of an equal volume of water. 8,6 grains of the compound, carefully freed from the saline solution in which it was produced, filled a space equal to that filled by 5,2 grains of water, consequently its specific gravity is 1,653.

When the compound is cooled artificially, either in water or in solution of nitrate of ammonia, the fluid surrounding it congeals at a temperature a little below 40° FAHRENHEIT, which seems to be owing to its becoming a solution of chlorine; for, as I have stated in a paper published in the Philosophical Transactions, the saturated solution of chlorine in water freezes very readily. The congelation of the fluid, in contact with the new compound, led me, when I first operated on it in very small quantities, to suppose it readily rendered solid by cooling; but I find in experimenting upon it, out of the contact of water, that it is not frozen by exposure to a mixture of ice and muriate of lime.

The compound gradually disappears in water, producing azote, and the water becomes acid, and has the taste and smell of a weak solution of nitro-muriatic acid.

The compound, when introduced into concentrated solution of muriatic acid, quickly resolves itself into gas, producing much more than its own weight of elastic fluid, which proves to be pure chlorine, and the solution evaporated affords muriate of ammonia.

In concentrated nitric acid it afforded azote.

In diluted sulphuric acid it yielded a mixture of azote and oxygen.

It detonated in strong solutions of ammonia. In weak solutions it produced azote.

It united to or dissolved in sulphurane, phosphorane, and alcohol of sulphur, without any violence of action, and dissolved in moderately strong solution of fluoric acid, giving it the power of acting upon silver.

When it was exposed to pure mercury, out of the contact of water, a white powder and azote were the results.

The first attempt that I made to determine the composition of the detonating substance, after my accident, was by raising it in vapour in exhausted vessels, and then decomposing it by heat; but in experiments of this kind, even though the whole of the substance was expanded into elastic matter, yet the vessel was often broken by the explosion, and in several instances violent detonations occurred during the process of exhaustion, probably from the contact of the vapour of the substance with the oil used in the pump.

In the only instance in which I was able to examine the products of the explosion of the substance in an exhausted vessel, no muriatic acid or water was formed, and chlorine and azote were produced; but it was impossible to form any correct opinion concerning the proportions of the gaseous matter evolved, as an unknown quantity of common air must have remained mixed with the vapour in the vessel.

The action of mercury on the compound appeared to offer a more correct and less dangerous mode of attempting its analysis; but on introducing two grains under a glass tube

filled with mercury and inverted, a violent detonation occurred, by which I was slightly wounded in the head and hands, and should have been severely wounded, had not my eyes and face been defended by a plate of glass attached to a proper cap, a precaution very necessary in all investigations of this body.

In using smaller quantities and recently distilled mercury, I obtained the results of the experiments, without any violence of action; and though it is probable that some accidental circumstance might have occasioned the explosion of the two grains, yet I thought it prudent, in my subsequent experiments, to employ quantities which, in case of detonation, would be insufficient to do any serious mischief.

In the most accurate experiment that I made, $\frac{1}{10}$ths of a grain of the compound produced, by its action upon mercury, 49 grain measures of azote. I collected the white powder which had been formed in this and other operations of the same kind, and exposed it to heat. It sublimed unaltered, without giving off any elastic or fluid matter, which there is the greatest reason to believe would not have happened, if the compound had contained hydrogen, or oxygen, or both. The sublimed substance had the properties of a mixture of corrosive sublimate and calomel.

If the results of this experiment be calculated upon, it must be concluded that the compound consists of 57 of azote to 648 of chlorine in weight, or 19 to 81 in volume; but this quantity of azote is probably less than the true proportion, as there must have been some loss from evaporation, during the time the compound was transferred, and it is possible that a minute quantity of it may have adhered to mercury not immediately within the tube.

The decomposition in this process is very simple, and must be supposed to depend merely upon the attraction of the mercury for chlorine, in consequence of which the azote is set free; and if the result does not strictly demonstrate the proportions of chlorine and azote in the compound, yet it seems at least to shew, that these are its only constituents.

As muriate of ammonia and chlorine are the only products resulting from its action upon solution of muriatic acid, it seems reasonable to infer, that this action depends on a decomposition of part of the muriatic acid, by the attraction of the new compound for hydrogen to form ammonia, which, at the moment of its production, combines with another portion of the acid, the chlorine of both compounds being set free.

On this view, the quantity of chlorine formed from a certain quantity of the compound being known, it becomes easy to determine the composition of the compound; for, ammonia being formed of three volumes of hydrogen and one of azote, and muriatic acid of one volume of hydrogen and one of chlorine, it is evident, that for every three volumes of chlorine evolved by the decomposition of muriatic acid, one volume of azote must be detached from the compound; and the weight of chlorine in the compound must be less than the weight of the whole quantity of chlorine produced by a portion, which is to the azote in the compound as 295 to 2295, if the relative specific gravities of the two gases be considered as 2,627 and 1.

Two grains of the compound, when exposed at the temperature of 62° FAHRENHEIT, and under a pressure of the atmosphere equal to that of 30,1 inches of mercury to strong solution of muriatic acid in a proper apparatus, afforded 3,91 cubic inches of chlorine.

In another experiment, one grain of the compound afforded 1,695 cubic inches of chlorine.

In a third experiment, one grain produced only 1,59 cubic inches.

In the two last experiments the compound was acted upon much more slowly, and the gas generated exposed to a much larger surface of solution of muriatic acid, and the appearance of a smaller relative proportion of chlorine must be ascribed to the absorption of a larger proportion of that gas by the liquid acid; and I found by exposing concentrated solution of muriatic acid to chlorine, that it soon absorbed nearly its volume of that gas.

I attempted to remove the source of error in the experiment, by using liquid muriatic acid holding chlorine in solution; but in this case the quickness of the action of the compound on the acid was greatly diminished, and it not being easy to obtain the point of absolute saturation of the acid with chlorine, some of the gas was absorbed in the nascent state during its slow production; and in most of my experiments made in this manner, I obtained less chlorine from a given weight of the compound, than in operating on pure solution of muriatic acid.

Liquid muriatic acid, whether concentrated or diluted in its pure state, does not affect the colour of the sulphuric solution of indigo; but it is immediately destroyed by solutions containing chlorine dissolved in them. The quantity of solution of indigo, which is deprived of colour by a given quantity of solution of chlorine, is directly as the proportion of chlorine it contains; and I found that the same quantity of chlorine, whether dissolved in a large or a small quantity of solution of

muriatic acid, destroyed the colour of the same quantity of the blue liquor.

On this circumstance it was easy to found a method of determining the precise quantity of chlorine produced in solution of muriatic acid, from a given quantity of the compound; namely, by comparing the power of a given quantity of muriatic acid, containing a known quantity of chlorine, to destroy the colour of solutions of indigo, with that of the muriatic acid, in which the compound had produced chlorine.

Two experiments were made. In the first, a grain of the compound was exposed on a large surface beneath a tube inverted in about six cubic inches of solution of muriatic acid, and the chlorine absorbed by agitation as it was formed. The acid so treated destroyed the colour of seven cubic inches of a diluted sulphuric solution of indigo; and it was found, by several comparative trials, that exactly the same effect was produced in another equal portion of the same solution of indigo, by 2,2 cubic inches of chlorine dissolved in the same quantity of muriatic acid.

In the second experiment, 1,8 cubic inches of chlorine were evolved in the gaseous form, the thermometer being at 58°, and barometer at 30,33, and suffered to pass into the atmosphere; and by the test of the solution of indigo, it was found that ,75 of a cubic inch remained dissolved in the acid.

Now, if the mean of these two experiments be taken, it appears that 1,61 grains of chlorine are produced in solution of muriatic acid by the action of a grain of the compound; and calculating on the data just now referred to, the compound must consist of 91 of chlorine and 9 of azote in weight, which

in volume will be nearly 119 to 30; and this estimation differs as little as might be expected from that gained by the action of mercury upon the compound.

It may fairly be concluded, that M. Gay Lussac's principle of the combination of gaseous bodies, in definite volumes, strictly applies to this compound, and that it really consists of four volumes of chlorine to one of azote; and the volumes likewise exactly coincide with the laws of definite proportions; and the detonating compound may be regarded as composed of one proportion of azote 26, and four proportions of chlorine 261.

I attempted a comparative experiment on the proportions in the compound, by estimating the quantity of azote produced in the decomposition of ammonia by it; but I found that this process was of no value for the purpose of analysis, for water appeared to be decomposed at the same time with the ammonia, and nitric acid formed; and, in consequence, the quantity of azote evolved was much less than it would have been, supposing the ammonia decomposed by the mere attraction of chlorine for hydrogen.

The results of the analysis of the new compound are interesting for several reasons.

They shew, what seemed probable from other facts, that there is no strict law of analogy, which regulates the combinations of the same substance with different substances. As three of hydrogen combine with one of azote, and one of hydrogen with one of chlorine, I thought it probable that the new compound would contain three of chlorine to one of azote, which is not the case.

This compound is the first instance known of one proportion

of a substance uniting to four proportions of another substance, without some intermediate compound of 1 and 1, 1 and 2, and 1 and 3; and the fact should render us cautious in adopting hypothetical views of the composition of bodies from the relations of the quantities in which they combine. Those who argue that there must be one proportion of oxygen in azote, because there ought to be six proportions in nitric acid, instead of five, which are produced from it by analysis, might with full as much propriety contend, that there must be azote in chlorine in some simple multiple of that existing in the compound.

It may be useful to shew, that many hypotheses may be framed upon the same principles; and which, consequently, must be equally uncertain. Views of this nature may be important in directing the practical chemist in his researches; but the philosopher should carefully avoid the developement of them with confidence, and the confounding them with practical results.

The compound of chlorine and azote agrees with the compounds of the same substance with sulphur, phosphorus, and the metals, in being a non-conductor of electricity; and these compounds are likewise decomposable by heat, though they require that of Voltaic electricity.

Sulphur combines only in one proportion with chlorine; and hence the action of *Sulphurane*, or Dr. THOMSON's muriatic liquor upon water, like that of the new compound, is not a simple phenomenon of double decomposition.

It seems proper to designate this new body by some name: *Azotane* is the term that would be applied to it, according to my ideas of its analogy to the other bodies which contain chlorine; but I am not desirous, in the present imperfect and

fluctuating state of chemical nomenclature, to press the adoption of any new word, particularly as applied to a substance not discovered by myself.

I am, my dear Sir,

very sincerely yours,

HUMPHRY DAVY.

XXIX. *Experiments on the Production of Cold by the Evaporation of the Sulphuret of Carbon.* *By* Alexander Marcet, *M. D. F. R. S. one of the Physicians to Guy's Hospital.*

Read July 8, 1813.

I HAD the honour, at an earlier period of this session, of giving to the Society, conjointly with Professor BERZELIUS, an account of the alcohol of sulphur, or sulphuret of carbon, and of noticing the remarkable volatility of that fluid. I have, since that period, tried a variety of experiments on the subject, and having found this compound more volatile than any other known body, and capable of producing, by its evaporation, a degree of cold of proportional intensity, I have been induced, in order to render the history of the sulphuret of carbon more complete, to present to the Society a brief account of my experiments.*

If the bulb of a small spirit thermometer be closely covered with a bag of fine flannel, or still better, with a piece of fine

* I would recommend to those, who may wish to repeat these experiments, to prepare this substance by means of a large earthen tube of about one inch and a half in diameter, instead of the small porcelain tubes, which are commonly used for this purpose. The process is always a tedious one; but a much more considerable quantity of the sulphureous liquor is procured by the larger tube in an equal space of time. The same tube can scarcely ever be used twice. About half a pint of the liquor may be obtained in one process; but the operation requires almost a whole day. Fresh pieces of sulphur may be successively introduced whilst the distillation is going on, and without renovating the charcoal, as the occasional admission of air does not materially affect the operation. The process would, no doubt, admit of further improvement.

lint, and a few drops of the sulphureous liquor poured upon it, the thermometer rapidly sinks from 60° to about 0; whilst if a similar experiment be made with ether, or alcohol, under the same circumstances, the temperature is reduced by the ether to only about + 20°, and by the alcohol to about 50°.* The hygrometrical state of the atmosphere appears to have some influence on these results.

If the bulb of the thermometer, thus wetted with the sulphureous liquor, be introduced into the receiver of an air-pump, (by means of a brass plate, as expressed in the annexed



sketch, the plate fitting the receiver air tight when laid upon its open neck), the thermometer sinks rapidly, as the ex-

* I have also tried some experiments on the comparative elastic force of the vapour of these fluids, by introducing in succession a few drops of them, through mercury, into a torricellian tube; and I obtained the following results:

	Sp. Grav.	*Temper.*	*Depression of Mercury.*	*Cold produced by Evaporation.*
Pure alcohol	0,806	66°	1,65	from 68° to 52°.
Sulphuret of carbon	1,272	66°	10,75	from 68° to 10°.
Rectified ether	0,724	66°	15,65	from 68° to 24.°

It would appear, therefore, that the degree of volatility of a fluid, or of cold produced by its evaporation, are far from bearing an exact proportion to the elastic force of its

haustion proceeds, and in one or two minutes it descends from + 70° to — 70°, or even 80° of FAHRENHEIT's scale, that is at least 40 degrees below the freezing point of mercury. A cold of — 65 or — 70 is easily obtained by exhausting the receiver till the rarefied air supports only one quarter of an inch of mercury; but with a pump capable of exhausting to one-eighth of an inch, the thermometer sinks to — 81° or 82° in less than two minutes. It is scarcely necessary to observe that, if instead of a spirit thermometer, a tube containing mercury be employed, this metal may be instantly frozen, and the remarkable contraction, which it undergoes at the moment of its congelation, may be seen and estimated with great facility.*

These experiments are but slightly influenced either by the temperature of the atmosphere, or by that of the sulphureous liquor; but if the air be very damp, the moisture which accumulates upon the bulb whilst cooling, slightly impedes the process. This moisture appears in the form of a hoar frost, forming a snowy arborescent deposition all over the covering of the bulb. The oily liquor itself does not freeze when exposed to a temperature of at least — 60°.

I shall not take up the time of the Society by a detail of the various attempts which I have made to increase still farther

vapour. It may be observed also, that the elastic force of the vapour of alcohol of sulphur was stated by Mr. BERZELIUS and myself, in our joint paper, to be only 7,36 inches, instead of 10,75; but this difference is accounted for from the circumstance of the temperature, which in that instance was 53,5°, instead of 66°; a difference which is very nearly that which might have been expected from a comparison with the force of ether at the same temperature as given in Mr. DALTON's tables.

* Mercury may also be frozen by a similar process, by means of ether, as I have stated in NICHOLSON's Journal for February last.

the effects just related. The fact is, that all these attempts have hitherto failed, and the simplest mode of performing the experiment has uniformly yielded the best results. Indeed, I am now inclined to believe, that unless a substance were found which should exert an attraction upon the sulphureous liquor, without being itself evaporable, (as is the case with the sulphuric acid, in Mr. LESLIE's mode of freezing water), I have nearly reached the limit of cold that can be produced in this manner. It may be proper to add, that the presence of sulphuric acid in this experiment would have no immediate effect, as it exerts no attraction upon the volatilized liquor; yet by drying the rarefied atmosphere which surrounds the bulb, it might perhaps assist a little in promoting the process of evaporation. It may also be useful to caution those who may wish to repeat these experiments, against working the pump too rapidly; for in this case, a degree of heat is produced (in consequence, no doubt, of the condensed air rushing violently through the valve), which is sufficient to inflame the volatilized liquor mixed with it. This happens most frequently with pumps of large dimensions, and is immediately perceived by the sulphureous fumes which issue from the barrel; no other consequence ensues, however, except the injury done to the valves, which, at all events, are apt to be acted upon, and more or less deranged, by the repeated contact of the volatilized liquor. On this account the receiver used for these experiments should be of the capacity of two or three pints; for if it be very small, and the vapour very concentrated, the valves appear to be proportionally injured.

XXX. *On a saline Substance from Mount Vesuvius.* *By* James Smithson, *Esq. F. R. S.*

Read July 8, 1813.

It has very long appeared to me, that when the earth is considered with attention, innumerable circumstances are perceived, which cannot but lead to the belief, that it has once been in a state of general conflagration. The existence in the skies of planetary bodies, which seem to be actually burning, and the appearances of original fire discernible on our globe, I have conceived to be mutually corroborative of each other; and at the time when no answers could be given to the most essential objections to the hypothesis, the mass of facts in favour of it fully justified, I thought, the inference that our habitation is an extinct comet or sun.

The mighty difficulties which formerly assailed this opinion, great modern discoveries have dissipated. Acquainted now, that the bases of alkalies and earths are metals, eminently oxydable, we are no longer embarrassed either for the pabulum of the inflammation, or to account for the products of it.

In the primitive strata, we behold the result of the combustion. In them we see the oxyd collected on the surface of the calcining mass, first melted by the heat, then by its increase arresting farther combination, and extinguishing the fires which had generated it, and in fine become solid and crystallized over the metallic ball.

Every thing tells that a large body of combustible matter still remains enclosed within this stony envelope, and of which volcanic eruptions are partial and small accensions.

Under this point of view, an high interest attaches itself to volcanoes, and their ejections. They cease to be local phenomena; they become principal elements in the history of our globe; they connect its present with its former condition; and we have good grounds for supposing, that in their flames are to be read its future destinies.

In support of the igneous origin, here attributed to the primitive strata, I will observe, that not only no crystal imbedded in them, such as quartz, garnet, tourmaline, &c. has ever been seen enclosing drops of water; but that none of the materials of these strata contain water in any state.

a. The present saline substance was sent to me from Naples to Florence, where I was, in May 1794, with a request to ascertain its nature. The general examination which I then made of it, shewed it to be principally what was at that time called *vitriolated tartar*, and it was in consequence mentioned as such in an Italian publication soon after. But as this denomination, surprising at that period, was not supported by the relation of any experiments, or the citation of any authority, no attention was paid to it; and the existence of this species of salt, native in the earth, has not been admitted by mineralogists, no mention being made of it, I believe, in any mineralogical work published since.

b. I was informed by letter, that it had "flowed out liquid from a small aperture in the cone of Vesuvius," and which I apprehend to have happened in 1792 or 1793.

c. The masses of this salt are perfectly irregular, their texture compact, their colour a clouded mixture of white, of the green of copper, and of a rusty yellow, and in some places are specks and streaks of black.

d. A fragment melted on the charcoal at the blow-pipe formed hepar sulphuris.

e. A piece weighing 9,5 grains was so strongly heated in a platina crucible, that it melted and flowed level over the bottom of it, but did not lose the least weight.

f. Not the slightest fume could be perceived on holding a glass tube wetted with marine acid over some of this salt, while triturating in a mortar with liquid potash; but a similar mixture being made in a bottle, and which was immediately closed with a cork, to which was fixed a bit of reddened litmus paper, the blue colour of the paper was restored.

g. Being dissolved in water, there was a small sandy residue, which consisted of green particles of a cupreous nature, of a yellow ochraceous powder, and of minute crystals of a metallic aspect of red oxyd of iron, by which the black spots in the mass had been occasioned.* Mr. KLAPROTH found a similar admixture in muriate of soda from Vesuvius.†

h. The solution had a feeble green tint. It did not alter blue or reddened turnsol paper.

i. Prussiate of soda-and-iron threw down a small quantity of red prussiate of copper from it. Liver of sulphur and tincture of galls likewise caused very small precipitations.

* What mineralogists denominate speculary iron ore, *Fer oligiste* of Mr. HAÜY, appears to be merely red oxyd of iron in crystals; red hematite the same substance in the state of stalactite; and red ochres the same in a pulverulent form. The hematites which afford a yellow powder are hydrates of iron.

† Essays, Vol. II. p. 67, Eng. Trans.

j. Carbonate of soda, and oxalate of potash, and solutions of magnesia, clay, copper, iron, and zinc, either had no effects, or extremely slight ones.

k. Solution of sulphate of silver produced a white curd-like precipitate. 9,85 grains of this salt (the weight of the insoluble matter being deducted) afforded 1,05 grains of slightly melted muriate, or chloride, of silver. This precipitate was equally produced after the salt had been made strongly red hot, so that it was not owing to a portion of sal ammoniac.

l. Tartaric acid, and muriate of platinum, occasioned the precipitates in its solution which indicate potash.

m. Nitrate of lime did not form any immediate precipitate in a dilute solution of it; but in a short time, numerous minute prismatic crystals of hydrate of sulphate of lime were generated.

n. Nitrate of barytes poured into a solution containing 9,8 grains of this salt afforded a precipitate, which after being ignited weighed 12,3 grains. The filtered solution crystallized entirely into nitrate of potash mixed with a few rhomboides of nitrate of soda.

o. Some of this salt finely pulverized was treated with alcohol. This alcohol on exhaling left a number of minute cubic crystals, which proved, by the test of nitric acid, to be muriate of soda. Prussiate of soda-and-iron caused a red precipitate of prussiate of copper in this alcoholic solution.

p. The solution of this salt afforded, by crystallization, sulphate of potash in its usual forms, and some prismatic crystals of hydrate of sulphate of soda.

q. To discover what had occasioned the precipitate with galls, (*i*) since copper has not this quality, a portion of this

salt, which had been recovered by evaporation from a filtered solution of it, was made red hot in a platina crucible. On extraction of the saline part by water, a very small quantity of a black powder was obtained. Ammonia dissolved only part of it, which was copper. The rest being digested with muriatic acid, and prussiate of soda-and-iron added, a fine Prussian blue was formed.

r. From several of the foregoing experiments, it appeared that no sensible quantity of any of the mineral acids, besides the sulphuric and muriatic, existed in combination with alkali in this volcanic salt. But Mr. TENNANT, whose many and highly important discoveries have so greatly contributed to the progress of chemical science, having detected disengaged boracic acid amongst the volcanic productions of the Lipari islands, and suggested that it might be a more general product of volcanoes than had been suspected,* it became important to ascertain whether the presence of any in this salt proved Vesuvius likewise to be a source of this acid. Alcohol heated on a portion of it in fine powder, and then burned on it, did not however shew the least green hue in its flame.

s. To ascertain the proportions of the ingredients of this saline substance, the following experiments were made:

10 grains of sulphate of potash of the shops were dissolved in 200 grains of water, and an excess of muriate of platina added. The precipitate edulcorated with 100 grains of water, and dried on a water bath, weighed 24,1 grains.

10 grains of the saline part of the native salt, treated precisely in every respect in the same way, afforded 17,2 grains of precipitated muriate of platina-and-potash.

* Trans. of the Geolog. Soc.

If 24,1 grains of this precipitate correspond to 10 grains of sulphate of potash, 17,2 grains of it correspond to 7,14 grains of this salt.

It has been seen (*n*) that 10 grains of the saline part of this volcanic salt would have afforded 12,55 grains of sulphate of barytes.

But 7,14 grains of sulphate of potash form only 9,42 grains of sulphate of barytes,* and therefore the remaining 3,13 grains of sulphate of barytes would be produced by the sulphate of soda, and correspond to 1,86 grains of it in an arid state, or uncombined with ice.†

10 grains of the saline part of this native salt would have produced 1,12 grains of ignited muriate of silver (*k*). By accurate experiments 241 grains of ignited muriate of silver have been found to correspond to 100 grains of ignited muriate of soda.‡

Consequently the soluble portion of the present Vesuvian salt consists of

Sulphate of potash	- -	7,14
Sulphate of soda	- -	1,86
Muriate of soda	- -	0,46
Muriate of ammonia, Muriate of copper, Muriate of iron	- -	0,54
		10,00

t. The insoluble sandy residue (*g*) having been thoroughly

* Dr. Marcet on Dropsical Fluids.

† Prof. Klaproth's Essays, Vol. I. p. 282.

‡ Dr. Henry, Phil. Trans. 1810.

edulcorated, dilute nitric acid was put to it. A green solution formed without any effervescence. Acetate of barytes scarcely rendered this solution turbid; but nitrate of silver produced a copious curd-like precipitate, and iron abundantly threw down copper from it. The green grains enclosed in this native sulphate of potash, appear, therefore, to be a submuriate of copper, of the same species as that of the green sands of Peru and Chili.

Muriatic acid dissolved the yellow ochraceous powder, and prussiate of soda-and-iron produced Prussian blue. I am inclined to believe this yellow powder to be a submuriate of iron, but its small quantity, and the admixture of the submuriate of copper, were impediments to entirely satisfactory results. Such a submuriate of iron, though, if I mistake not, overlooked by chemists, exists, for the precipitate which oxygen occasions in solution of green muriate of iron, contains marine acid.

Possibly this yellow powder, and the crystals of speculary iron which exist in this Vesuvian salt, have been produced by a natural sublimation of muriate of iron, similar to that of the experiment of the Duke d'AYEN, recorded by MACQUER,* and which was known long before to Mr. BOYLE and Dr. LEWIS.†

This Vesuvian salt, considered in its totality, has presented no less than nine distinct species of matters, and a more rigorous investigation, than I was willing to bestow on it, would probably add to their number.

July 3, 1813.

* *Dict. de Chemie, Art. Fer.*

† A course of practical chemistry by WILLIAM LEWIS, 1746, page 63, note *f*.

XXXI. *Some Experiments and Observations on the Substances produced in different chemical Processes on Fluor Spar. By Sir* Humphry Davy, *LL.D. F.R.S. V.P.R.I.*

Read July 8, 1813.

In the Bakerian Lecture, for 1808, I have given an account of an experiment on the combustion of potassium in silicated fluoric acid gas, in which the gas was absorbed, and a fawn coloured substance formed, which effervesced with water, and left, after its action on that fluid, a residuum which burnt when heated in oxygen, reproducing silicated fluoric acid gas; and I concluded from the phenomena, that the acid gas was decomposed in the process, that oxygen was probably separated from it by the potassium, and that the combustible substance was a compound of the siliceous and fluoric bases.

The experiment of burning potassium in silicated fluoric acid gas was made likewise by M. M. Gay Lussac and Thenard, before I published any account of my researches on this phenomenon. It was indeed one of the most obvious applications of potassium, and it occurred to many others, as well as to myself, that it might be made, immediately after I discovered that metal.

M. M. Gay Lussac and Thenard drew the same conclusions as I did, namely, that the acid gas was probably decomposed during the action of potassium on silicated fluoric acid;

but their general views differed from mine in this respect, as they supposed, that no part of the inflammable matter was derived from silica, and they likewise reasoned on the phenomena with more caution.

At the time that my conclusions were drawn, I was ignorant of the true nature of the muriatic acid. After I had tried in vain to decompose oxymuriatic gas, and after I had found that the compounds of this substance with phosphorus, sulphur, and the metals combined with ammonia without any decomposition, and produced compounds in which no oxygen could be discovered; I was forcibly struck by the analogy between the oxymuriatic and the fluoric compounds, and led to doubt of the justness of my ideas respecting the nature of fluoric acid.

I tried an experiment on the comparative quantities of fluate of lime, formed from equal volumes of silicated fluoric acid gas, one of which had been acted upon by potassium, and then exposed to solution of ammonia, the other had been absorbed by solution of ammonia: and I found the proportion of calcareous fluate nearly one-third larger in the latter case. This result at first seemed favourable to my early ideas, that the acid contained a peculiar inflammable basis, which was separated by the potassium, and existed in the combustible substance insoluble in water; but it could not be considered as decisive on the question, for, it occurred to me as possible, that this substance might be silicum, or the basis of silica united to a much smaller proportion of the fluoric principle, than that existing in silicated fluoric acid.

During the period that I was engaged in these investigations, I received two letters from M. AMPERE, of Paris, con-

taining many ingenious and original arguments in favour of the analogy between the muriatic and fluoric compounds. M. AMPERE communicated his views to me in the most liberal manner; they were formed in consequence of my ideas on chlorine, and supported by reasonings drawn from the experiments of M. M. GAY LUSSAC and THENARD.

Before I enter upon the detail of the investigations which promise to elucidate the nature of the fluoric compounds, it will be right to describe those substances produced from fluor spar, which have been the principal objects of my experiments, and to mention the different hypothetical views that may be formed respecting them.

The first of these substances is the silicated fluoric acid gas, which was discovered by SCHEELE, and examined in its pure state by PRIESTLEY. It is formed by heating a mixture of fluor spar, powdered glass, and sulphuric acid. It is a very heavy elastic fluid, its specific gravity being nearly forty-eight times as great as that of hydrogen. It produces, according to my brother Mr. JOHN DAVY, a quantity of silica equal to $\frac{153}{1000}$ of its own weight by its action upon water, and a quantity equal to $\frac{614}{1000}$ of its weight by its action upon solution of ammonia. It condenses twice its own volume of ammonia, and forms a solid salt, volatile when free from water without decomposition.

Liquid fluoric acid, the second of these substances, was discovered by SCHEELE, but first obtained in its pure form by M. M. GAY LUSSAC and THENARD. It is procured by heating concentrated sulphuric acid and pure fluor spar, in retorts of silver or lead, and receiving the product in receivers of the same metals artificially cooled. It is a very active substance, and

must be examined with great caution. According to my experiments, its specific gravity is 1,0609.* It produces a high degree of heat when mixed with water, and such is its degree of attraction for water, that it becomes denser by combining with that fluid. By adding water, in very small quantities at a time, to pure liquid fluoric acid, I found that its specific gravity gradually increased till it became 1,25: it is, I believe, the only known body possessed of this property.

The third substance is fluo-boric acid gas, which was discovered by M. M. GAY LUSSAC and THENARD. It is produced by intensely heating, in an iron tube, a mixture of dry boracic acid and fluor spar, or by gently heating in a glass retort a similar mixture with sulphuric acid. Its specific gravity is rather more than thirty-two times as great as that of hydrogen. It forms a solid salt, volatile without decomposition, by condensing its own volume of ammonia. The ammoniacal salt dissolved in water and distilled, affords boracic acid.

The most important phenomena of chemical change, in which these bodies operate, that may be supposed to illustrate their nature, is their agency upon potassium and other metals. The action of potassium upon silicated fluoric gas has been already referred to. M. M. GAY LUSSAC and THENARD, by heating potassium and sodium in fluo-boric acid gas, obtained fluate of potassa or soda, and the basis of the boracic acid; and by exposing potassium to liquid fluoric acid, their results were hydrogen and acid fluate of potassa.

* Unless it is distilled through tubes and into vessels of pure silver, its specific gravity is greater; it readily dissolves tin, and slowly dissolves lead, and after being long kept in vessels of pure silver, it is found to have taken up a small portion even of that metal.

Three hypotheses may, according to sound analogies, be formed on the nature of the fluoric combinations. In the first, which is that generally adopted, the silicated fluoric acid gas is supposed to be a compound of silica and a peculiar acid, itself consisting of inflammable matter and oxygen; fluo-boric acid gas, a compound of boracic acid and the same acid; and pure liquid fluoric acid as water combined with the acid.

In the second hypothesis, that which I have alluded to in the beginning of this paper, and that adopted by M. AMPERE, the silicated fluoric acid is conceived to consist of a peculiar undecompounded principle, analogous to chlorine and oxygen, united to the basis of silica, or *silicum*; the fluo-boric acid of the same principle united to boron; and the pure liquid fluoric acid as this principle united to hydrogen.

In the third hypothesis, which probably would have been formed by the disciples of the phlogistic school of chemistry, had they been acquainted with the facts, the liquid fluoric acid is considered as an undecompounded body; and the metals and inflammable bodies as compounds of certain unknown bases with hydrogen: silicated fluoric acid gas, on this idea, must be regarded as a compound of the fluoric acid with the basis of silicum, and fluo-boric acid gas as a compound of fluoric acid and the basis of boron.

Whoever will consider, with attention, the different facts that have been brought forward by SCHEELE, GAY LUSSAC and THENARD, JOHN DAVY, and myself, will find that they will admit of explanation on either of these hypotheses; and, as in all the cases yet brought forward, of the most simple chemical action of other bodies on the fluoric substances, more than one new form of matter is produced, no explanation of

the phenomena can at present be given without involving suppositions.

It is not easy to devise simple experiments to ascertain which of these hypotheses is true, yet, in admitting strict analogical reasoning, it is easy to shew which is most conformable to the general series of chemical facts.

Those acids which are known by direct experiments of decomposition by heat, to consist of oxygen, bases, and water, such as the strongest sulphuric and nitric acids and hydrophosphorous acid, when they are acted on by ammonia, afford moisture: this is easily proved, by causing them to absorb ammoniacal gas in glass retorts, and gently heating the mixture, when water immediately appears. On this view, it occurred to me, if the liquid fluoric acid was a compound of water, and inflammable bases, and oxygen, that water ought to be produced when it was made to combine with ammonia. It was not possible to make the experiment in glass vessels, as the acid acts with great violence on glass, producing silicated fluoric acid gas. I had recourse, therefore, to an apparatus made of platina. A small tray of platina was filled with pure liquid fluoric acid, and introduced into a tube of platina connected by proper stop-cocks with a mercurial gazometer, filled with ammonia; the end of the platina tube was closed by a brass stopper, and a communication made between the ammonia and the fluoric acid; the ammonia was gradually absorbed, producing heat; and white fumes sometimes rose into the gas-holder, so that it was necessary from time to time to cut off the communication; ammoniacal gas was supplied till no more absorption took place. When the tube was quite cool, the stopper was removed, and the result examined; the

interior contained a white crystalline mass, but there was no appearance of fluid.* A polished brass tube, cooled by means of ice, was held over the aperture of the platina tube, and it was gently heated till the salt began to sublime, but no moisture was found condensed in the cold tube of brass.

This experiment is unfavourable to the idea, that the liquid fluoric acid contains water; and the following result is likewise unfavourable to the idea that it consists of an inflammable basis united to oxygen. Solid and perfectly dry fluate of ammonia was introduced into a tray of platina, with about an equal quantity of potassium, and the tray was heated in a small tube of glass connected with a mercurial apparatus. A violent action took place, gas was disengaged with great violence, which remained for some time clouded; the application of heat was continued till the tube was red: it was then suffered to cool, and the results examined. Much white matter, which proved to be fluate of potassa, had been carried by the violence of the action out of the tray of platina into the glass tube; and a little potassium had sublimed in the tube. The tray contained a considerable portion of potassium, and a saline matter, which had all the characters of fluate of potassa. The gas disengaged, consisted of ammonia and hydrogen, to each other in volume nearly as two to one; but the experiment cannot be considered as decisive on this point, as no particular precautions had been taken to dry the mercury.

* It is necessary that pure liquid fluoric acid, *i. e.* that which has the lowest specific gravity, be used for this experiment. The first time that I made it, I obtained moisture, owing to my having formed the hydro-fluoric acid by means of sulphuric acid that had not been previously boiled, and which must have contained more than one proportion of water.

Now, if there had existed oxygen combined with an inflammable basis in the fluate of ammonia, it might have been expected to have been separated, or at least to have formed a new combination during the action of potassium upon the fluate of ammonia, which is the case with such ammoniacal salts as contain acids in which oxygen is an element. Thus nitrate of ammonia acted on by potassium, as I have found, affords azote and ammonia; and sulphur is partly disengaged, and partly newly combined during the agency of potassium in excess upon sulphate of ammonia.

The action of potassium upon fluate of ammonia is precisely similar to its action upon muriate of ammonia, in which as I have found, by numerous experiments, ammonia and hydrogen to each other in volume as two to one are disengaged, and muriate of potassa (*potassane*) formed.

All the hydrates, that is, all the substances which contain definite proportions of water, united to acids, alkalies, or oxides, which are fluid, or capable of being rendered fluid by heat, when exposed to the chemical agency of Voltaic electricity, undergo decomposition, and their inflammable principles, either pure or combined with a smaller proportion of oxygen, are disengaged at the negative surface in the circuit, and their oxygen at the positive surface. Thus sulphuric acid affords sulphur and hydrogen at the negative surface, and the hydrophosphorous acid, phosphuretted hydrogen and phosphorus, and nitric acid nitrous gas; and all these bodies yield oxygen at the positive surface.

I undertook the experiment of electrizing pure liquid fluoric acid, with considerable interest, as it seemed to offer the most probable method of ascertaining its real nature; but consider-

able difficulties occurred in executing the process. The liquid fluoric acid immediately destroys glass, and all animal and vegetable substances; it acts on all bodies containing metallic oxides; and I know of no substances which are not rapidly dissolved or decomposed by it, except metals, charcoal, phosphorus, sulphur, and certain combinations of chlorine.

I attempted to make tubes of sulphur, of muriates of lead and of copper containing metallic wires, by which it might be electrized, but without success. I succeeded, however, in boring a piece of horn silver in such a manner, that I was able to cement a platina wire into it, by means of a spirit lamp, and by inverting this in a tray of platina filled with liquid fluoric acid, I contrived to submit the fluid to the agency of electricity in such a manner, that in successive experiments it was possible to collect any elastic fluid that might be produced. Operating in this way, with a very weak Voltaic power, and keeping the apparatus cool by a freezing mixture, I ascertained that the platina wire at the positive pole rapidly corroded, and became covered with a chocolate powder; gaseous matter separated at the negative pole, which I could never obtain in sufficient quantities to analyze with accuracy; but it inflamed like hydrogen. No other inflammable matter was produced when the acid was pure.

In a case in which the acid had been condensed in a tube of lead, joined by a solder containing tin, a large quantity of powder separated at the negative surface of a dark colour, and which appeared to be tin mixed with a subfluate; the powder burnt when heated in the air, and gave fluoric fumes when treated by potassa and sulphuric acid.

I attempted to electrize the liquid fluoric acid, by making plumbago the positive surface; but the plumbago was quickly destroyed, a subfluate of iron was deposited on the negative surface, and the liquid became turbid and black. When a point of charcoal attached to a wire of platina was made positive, the effects were similar to those produced by a platina wire alone, for the acid speedily penetrated through the pores of charcoal, and the platina, in consequence, became a point of contact with the fluid.

I applied the power of the great Voltaic batteries of the Royal Institution to the liquid fluoric acid, so as to take sparks in it. In this case, gas appeared to be produced from both the negative and the positive surfaces; but it was probably only the undecompounded acid rendered gaseous, which was evolved at the positive surface, for during the operation the fluid became very hot, and speedily diminished. The manner in which the surrounding atmosphere became filled with the fumes of the fluoric acid, rendered it, indeed, very difficult to examine the results of any of these experiments; the dangerous action of these fumes have been described by M. M. GAY LUSSAC and THENARD, and I suffered considerable inconvenience from their effects during this investigation. By mere exposure to them in their uncondensed state, my fingers became sore beneath the nails, and they produced a most painful sensation, which lasted for some hours, when they came in contact with the eyes.

The phenomena of the Voltaic electrization of fluoric acid, present no evidences in favour of its containing a peculiar combustible substance and oxygen; and the most simple mode of explaining them, is by supposing the fluoric acid, like

muriatic acid, composed of hydrogen, and a substance, as yet unknown, in a separate form, possessed, like oxygen and chlorine, of the negative electrical energy, and hence determined to the positive surface, and strongly attracted by metallic substances.

This view is much more conformable to the general order of chemical and electrical facts than the third hypothesis, just now mentioned.

It is indeed possible to conceive, if the metals be regarded as compounds of hydrogen, that the hydrogen may be produced from the metal, positively electrified at the time that the acid combines with its supposed basis, and that this hydrogen may be transferred to the negative surface; but this supposition involves a multitude of others; and the results of the electrization of fluoric acid are analogous to most of the results of the electrization of water and muriatic acid, both of which are shewn by analysis and synthesis to be compounds of hydrogen; and in the electrical decomposition of these bodies, their characteristic element is generally combined with the positive metallic surface.

In the Bakerian Lecture for 1810, I have given an account of the action of potassium upon pure silica. In this process, the potassium acquires oxygen, and a combustible substance, which consists either of the basis of silica, or the basis of silica combined with potassium appears. In supposing the silicated fluoric acid gas to be composed of this basis and the fluoric principle, it is easy to explain the action of potassium upon it, and the complicated phenomena, occasioned by the agency of water, and acids, and oxygen, on the results of this action. The potassium must be conceived to attract a part of the fluoric

principle from the siliceous basis, or to form a triple compound, from which silicated fluoric acid gas is capable of being reproduced, in consequence of the combination of a part of the potassium and siliceous basis with oxygen; and on this idea the cause of the apparent loss of the fluoric principle, in the experiments on the action of ammonia on the product of the combustion of potassium in silicated fluoric acid gas, becomes obvious.

Assuming then from the analogy with chlorine, that the different fluoric compounds consist of inflammable bodies united to a peculiar principle, it follows that all attempts to decompose the fluoric acids, by combustible substances, can lead to no other result, than that of occasioning new combinations of the fluoric principle; and the only methods which seemed plausible for obtaining this principle pure, after that by electrical decomposition had failed, were by the action of oxygen or chlorine on certain of its compounds. Chlorine is, in certain instances, detached from hydrogen by oxygen; and oxygen, in a number of cases, is detached from metals by chlorine; I thought it therefore probable, that the fluoric principle might, in some process, be separated from bases by either chlorine or oxygen.

In selecting compounds for experiments of this kind, I was guided by the relative attractions of the fluoric and muriatic acids, of chlorine and oxygen. Horn silver and calomel, and muriate of potassa are not decomposed by fluoric acid, but fluate of silver, of mercury, and of potassa are easily decomposed by muriatic acid; I therefore conceived, that the fluoric principle would most likely be expelled from the dry fluates of silver, mercury, and potassa by chlorine.

I made some pure fluates of silver and mercury, by dissolving the oxides of these metals in fluoric acid, and I heated them in small trays of platina; much fluoric acid was driven off in this process, which I continued in the case of the fluate of mercury till the salt began to sublime, and in that of the fluate of silver till it was red hot.

The dry salts were introduced in small quantities into glass retorts, which were exhausted and then filled with pure chlorine: the part of the retort in contact with the salt was heated gradually till it became red. There was soon a strong action, the fluate of mercury was rapidly converted into corrosive sublimate, and the fluate of silver more slowly became horn silver. In both experiments there was a violent action upon the whole of the interior of the retort. On examining the results, it was found that in both instances there had been a considerable absorption of chlorine, and a production of silicated fluoric acid gas, and oxygen gas.

I tried similar experiments, with similar results, upon dry fluates of potassa and soda. By the action of a red heat, they were slowly converted into muriates with the absorption of chlorine, and the production of oxygen, and silicated fluoric acid gas, the retort being corroded even to its neck.

The obvious explanation of these phenomena is, that a particular principle, the acidifying matter of the fluoric acid, combined with the metals, is expelled from them by the stronger attraction of the chlorine, and that this principle coming in contact with glass decomposes it by its attraction for the silicum and sodium, and separates them from the oxygen with which they were combined.

I made various attempts to procure the fluoric principle in a

pure form. I heated the fluates of potassa and soda in trays of platina, in a tube of platina connected with a vessel filled with chlorine. In this case the fluates were converted into muriates, with a considerable increase of the weight of the tray; and the platina was violently acted upon, and covered with a reddish brown powder; and in the instance in which fluate of potassa was used, a compound of fluate of platina and muriate of potassa was formed.

There was a considerable absorption of chlorine; but no new gaseous matter could be discovered in the gas in the tube.

I tried to obtain the fluoric principle pure, by decomposing the fluates in a tube of silver, but with no better success; the silver was acted upon both by the chlorine and the fluoric principle, and rapidly dissolved. I used glass tubes coated with resin of copper (*cuprane*) and hornsilver (*argentane*), on which I concluded that the fluoric principle would have no action from the decomposition of fluate of silver by chlorine; but at the degree of heat required to decompose the fluoric salts, the muriates were always fused, the glass violently acted upon, and silicated fluoric acid gas formed.

In one instance, in which fluate of potassa had been heated in a platina tray and tube, in which muriate of potassa had been fused, for the purpose of defending the interior, as much as possible, from the action of the fluoric principle, the gas, when disengaged into the atmosphere, had a peculiar smell, different from that of chlorine, (which certainly formed the greatest proportion of the elastic matter,) and more disagreeable; and dense white fumes were produced by its action upon the air. A portion of this gas thrown into a glass receiver, over

mercury, acted upon the glass, and silicated fluoric acid gas was generated. On examining the platina tray, however, it was found corroded, and the reddish brown powder formed.

In the course of these investigations, I made several attempts to detach hydrogen from the liquid fluoric acid, by the agency of oxygen and chlorine. It was not decomposed when passed through a platina tube heated red with chlorine, nor by being distilled from salts containing abundance of oxygen, or those containing abundance of chlorine.

I distilled the fluates of lead and mercury with phosphorus and sulphur, with the hope of obtaining compounds of the fluoric principle with phosphorus and sulphur. In all experiments of this kind, a decomposition took place, and the glass tubes employed were violently acted upon, and sulphurets and phosphurets were formed. When I used tubes lined with sulphur the decomposition was less perfect; but minute quantities of limpid fluid condensed in a part of the tube cooled by ice, both in the cases when sulphur and when phosphorus were used; it had the appearance of hydrofluoric acid, and speedily dissipated itself in white fumes. Whether they were that substance which had obtained its hydrogen from these inflammable bodies, or compounds of sulphur and phosphorus with the fluoric principle, I have not ascertained, but the first opinion seems most probable.

When I heated fluate of lead and finely powdered charcoal strongly in the air, the lead became revived, and white fumes were produced. I thought it probable, that in this case a compound of fluorine and charcoal was formed; but on trying the experiment in a close vessel of platina, no change took

place; and it evidently depended upon the presence of hydrogen in the vapour of the atmosphere, or in the flame of the spirit lamp, by which the experiment was made, and I found muriate of silver decomposed, and silver produced under the same circumstances.

From the general tenor of the results that I have stated, it appears reasonable to conclude that there exists in the fluoric compounds a peculiar substance, possessed of strong attractions for metallic bodies and hydrogen, and which combined with certain inflammable bodies forms peculiar acids, and which, in consequence of its strong affinities and high decomposing agencies, it will be very difficult to examine in a pure form, and, for the sake of avoiding circumlocution, it may be denominated *fluorine*, a name suggested to me by M. AMPERE.

From experiments that I have made on the composition of the fluoric combinations, and which I shall soon have the honour of communicating to the Society, it appears that the number representing the definite proportion in which fluorine combines, is less than half the number representing that in which chlorine combines; and hydrates in becoming fluates lose weight, so that on the generally received idea of the existence of a peculiar acid in the fluates, and of their being compounds of oxides, with an acid containing oxygen, that acid, according to the law of definite proportions, must contain more oxygen in proportion to its quantity of inflammable matter than water, which is highly improbable, and contrary to all analogies.

Dr. WOLLASTON has found, that the fluoric combinations have very low powers of refracting light, and particularly the pure fluoric acid; so that the refracting powers of fluorine

will probably be found lower than those of any other substance, and it appears to possess higher acidifying and saturating powers than either oxygen or chlorine.

It is easy to perceive, in following the above theory, that all the ideas current in chemical authors respecting the fluoric combinations, must be changed. Fluor spar, and other analogous substances, for instance, must be regarded as binary compounds of metals and fluorine.

Many objects of enquiry arise, likewise, from these new views: the topaz contains the fluoric principle, but new experiments are required to shew whether that gem is a true silicated fluate of alumina, or a compound of the inflammable bases of alumina and silica with fluorine.

I have ascertained that the chryolite yields no silicated fluoric gas, when acted on by sulphuric acid, but merely pure fluoric acid; but I have not continued the research so far, as to determine whether it contains fluorine united to inflammable matter only, or fluorine and oxygen.

XXXII. *Catalogue of North Polar Distances of Eighty-four principal fixed Stars, deduced from Observations made with the Mural Circle at the Royal Observatory.* By John Pond, *Esq. Astronomer Royal, F. R. S.*

Read July 8, 1813.

THE Catalogue, which I have the honour to transmit to the Society, is deduced from the whole of the observations made with the mural circle, from its first erection in June, 1812, to the present time. I am still employed in endeavouring to give it a greater degree of precision, and when it is entirely finished, I propose to submit some of the observations themselves to the Society, and explain the method by which the results have been obtained.

I have already mentioned, that I use neither level nor plumb-line; but determine the position of the instrument by means of a standard catalogue of stars derived from the instrument itself, in such a manner, that every series of observation of these stars, serves the double purpose of ascertaining the position of the instrument, and at the same time improving the Catalogue.

As the present Catalogue has been formed by frequently changing the position of the telescope on the circle, for the purpose of correcting every possible error of division, my observations have not been calculated to decide the question of parallax which has been suspected to exist in α Lyræ, α

Aquilæ, and some other stars. But from this time forward, I propose to use the instrument in one position of the telescope, with the hope of ascertaining, if not the parallax of these stars, at least the limits, which it does not exceed.

Though the change of position in the telescope, by which all error of division is avoided, is one of the most beautiful properties of this instrument, yet so accurately is it divided, that I cannot perceive, with certainty, any effect produced by this change, for I have often found as great a discordance between two series of observations made on the same divisions, as when they are entirely changed by a new position of the telescope. What the error of division may amount to in any one position, I cannot exactly say; but, I think, when the six microscopes are used, it can never exceed half a second, and very rarely amounts to half that quantity.

That some opinion may be formed of the accuracy of this instrument, I have subjoined to the Catalogue the results of the observations of some of the standard stars, whose places I am anxious to determine with the greatest precision, since I propose in future to determine all north polar distances by comparison with these stars; precisely in the same manner as right ascensions are now determined by comparison with the thirty-six stars, whose places have been so accurately determined by Dr. MASKELYNE.

GENERAL CATALOGUE.

Names of Stars.	No. of Obs.	N. P. Distances Jan. 1, 1813.
		° ′ ″
1 γ PEGASI	25	75 51 21,0
2 α CASSIOP.	42	34 29 22,7
3 γ Cassiop.	8	80 17 53,0
4 POLARIS	200	1 41 21,75
5 δ Cassiop.	14	30 44 26,8
6 α ARIETIS	50	67 25 36.5
7 α CETI	18	86 39 0,75
8 α PERSEI	44	40 48 52,7
9 δ Persei	10	42 49 16,4
10 η Tauri	10	66 28 55,5
11 γ Eridani	9	104 2 51,2
12 γ Tauri	10	74 49 59,8
13 1 δ	9	72 54 18,3
14 2 δ	10	72 59 53,8
15 ε	10	71 14 39,2
16 ALDEBARAN	56	73 52 35,4
17 CAPELLA	80	44 12 20,5
18 RIGEL	30	98 25 33,8
19 β TAURI	50	61 33 43,6
20 γ Orionis	8	83 49 48,3
21 δ	18	90 26 48,6
22 ε	14	91 19 48,5
23 ζ	16	92 3 1,8
24 α ORIONIS	50	82 38 15,7
25 γ Geminorum	12	73 27 5,8
26 ε	11	64 41 41,4
27 SIRIUS	34	106 28 0,7
28 δ Geminorum	11	67 41 1,9
29 η Canis Major	6	118 56 42,7
30 CASTOR	30	57 42 46,7

Names of Stars.	No. of Obs.	N. P. Distances begin. 1813.
31 PROCYON	40	84° 18′ 14″,4
32 POLLUX	40	61 31 56,3
33 α HYDRÆ	10	97 51 11,3
34 ε Leonis	4	65 22 11,3
35 REGULUS	62	77 7 22,7
36 ζ Leonis	4	65 39 17,3
37 γ	4	69 12 59,7
38 α URS. MAJ.	60	27 14 31,5
39 δ Leonis	3	68 27 11,3
40 β LEONIS	24	74 22 57,7
41 γ URS. MAJ.	48	35 15 55,3
42 δ Urs. Maj.	13	31 55 38,2
43 κ Draconis	3	19 10 45,3
44 δ Virginis	3	85 34 58,6
45 α SPICA. VIRG.	20	100 10 51,3
46 η URS. MAJ.	80	39 44 57,9
47 α Draconis	12	24 43 38,2
48 ARCTURUS	80	69 50 19,1
49 θ Bootes	6	37 16 49,7
50 π	7	72 46 25,8
51 ι	12	62 7 52,4
52 1 } α LIBRÆ	15	105 15 22,7
53 2 }	4	105 12 38,7
54 β URS. MIN.	90	15 4 49,0
55 β Libra	8	98 41 3,4
56 α COR. BOR.	90	62 38 55,4
57 α SERPENTIS	70	82 38 39,3
58 δ Scorpii	9	112 4 41,4
59 1 β	8	109 16 55,7
60 2 β	7	109 16 43,6

Names of Stars.	No. of Obs.	N. P. Distances begin. 1813.
		° ′ ″
61 δ Ophiuchi	16	93 12 9,8
62 ANTARES	96	116 0 16,6
63 ζ Herculis	10	58 3 4,7
64 α HERCULIS	50	75 29 14,0
65 α OPHIUCHI	70	77 17 39,2
66 γ DRACONIS	90	38 29 3,7
67 α LYRÆ	100	51 23 0,5
68 ζ Aquilæ	13	76 24 19,0
69 δ Draconis	21	22 40 0,5
70 δ Aquilæ	10	87 14 53,4
71 γ Aquilæ	38	79 50 0,6
72 α AQUILÆ	100	81 36 58,7
73 β Aquilæ	12	84 3 4,1
74 1 α } CAPRICORNI	35	103 5 35,4
75 2 α }	28	103 6 52,3
76 α Delphini	10	74 44 28,9
77 α CYGNI	80	45 22 56,9
78 α CEPHEI	40	98 12 12,5
79 β Aquarii	13	96 23 11,9
80 β CEPHEI	50	20 5 30,6
81 δ Capricorni	12	106 58 6,6
82 α AQUARII	20	91 18 21,6
83 α PEGASI	20	75 47 51,8
84 α ANDROMEDÆ	35	61 56 29,6

RESULTS OF OBSERVATIONS.

β Ursæ Minoris.

1812, June 12, 13, 14, 16, 17, 21, 22, 23, 28, 29 -	49,659	Mean of 10	49,659
June 30, July 6, 8, 9, 18, 20, 21, 22, 28, Aug. 7 -	48,926	Mean of 20	49,292
Aug. 13, 15, 17, 18, 19, Sept. 15, 16, 20, 21, Oct. 2 -	48,996	Mean of 30	49,194
Oct. 4, 18, 13, 19, 21, 25, 26, 28, 29 - -	48,617	Mean of 40	49,049
Oct. 31, Nov. 9, 14, 18, 21, 22, 23, Dec. 7, 8, 12 -	48,485	Mean of 50	48,987
Dec. 14. 1813 May 21, 26, 27, 28, 29, 31, June 1, 4, 7	48,951	Mean of 60	48,989
June 8, 11, 12, 16, 21, 22, 25, July 4, 5, 6, - -	49,695	Mean of 70	49,047
July 9, 12, 16, 17, 18, 20, 22, 24, 25, 26 - -	48,651	Mean of 80	48,998
July 28, 29, 30, Aug. 4, 7, 9, 12, 13, 16, 18 -	48,616	Mean of 90	48,955

This result is probably exact to within a quarter of a second. The discordances seem quite accidental, they neither arise from parallax, nor error of division, for the three last series were made with the telescope in the same position, and consequently upon the same division, yet they differ more than observations usually do, which are made in different positions.

β Cephei.

1812, Oct. 22, 24, 26, 29, 31, Nov. 3, 4, 21, 22, Dec. 5	} 30″,574	Mean of 10	30″,574
Dec. 8, 9, 13. 1813, Mar. 16, 20, Apr. 1, 2, 3, 8, 14	} 30,348	Mean of 20	30,458
Aug. 11, 12, 13, 15, 16 17, 19, 20, 22, 23 - -	} 30,985	Mean of 30	30,634
Aug. 24, 25, 26, 30, 31, Sept. 2, 3, 4, 5, 6 - -	} 30,894	Mean of 40	30,699

This result is probably exact to a quarter of a second.

α Ursæ Majoris.

1812, June 13, 29, July 9, 10, 15, 20, Aug. 14, Oct. 1, 2, 3	} 31″,500	Mean of 10	31″,500
Oct. 4, 6, 14, 15, 18, 19, 20, 23, 27, 28 - -	} 31,596	Mean of 20	31,548
Oct. 30, Nov. 4, 5, 9. 1813, Mar. 23, 26, 29, 31, Apr. 2, 3	} 31,525	Mean of 30	31,540
Apr. 4, 7, 8, 9, 10, 12, 13, 14, 20, 26 - - -	} 31,731	Mean of 40	31,588
May 24, 26, 31, June 1, 7, 8, 9, 11, 16, 25 -	} 31,124	Mean of 50	31,495
June 26, 27, July 6, 10, 18, 20, 25, 30, Aug. 5, 6 -	} 31,279	Mean of 60	31,459

This result seems to be extremely exact, and probably does not differ more than one-tenth of a second from the truth.

α Cephei.

1812, Oct. 28, 29, 31, Nov. 3, 21, 22, Dec. 5, 8, 9, 13	} 12″,484	Mean of 10	12″,484
1813, Mar. 20, Apr. 1, 3, 8, 12, Aug. 11, 12, 13, 15, 16	} 12,735	Mean of 20	12,610
Aug. 17, 19, 20, 21, 22, 23, 24, 25, 30, 31 - -	} 12,435	Mean of 30	12,551
Sept. 2, 3, 4, 5, 6, 7, 8, 9, 10, 14 - -	} 12,228	Mean of 40	12,470

Exact to a quarter of a second.

α Cassiop.

1812, June 17, 22, 28, July 6, 7, Oct. 26, 28, 29, 31, Nov. 3	22,839	Mean of 10	22,839
Nov. 4, 5, 6, 7, 19, 20, 22, 28, 29, Dec. 6 - -	22,456	Mean of 20	22,647
Dec. 7, 8, 9, 10. 1813, Jan. 9, 10, 16, 22, 24, Apr. 8 -	522,59	Mean of 30	22,630
Apr. 9, 11, 17, May 26, 27, 28, 31, June 7, 10, 12 -	23,041	Mean of 40	22,726

This result may, I think, be relied on, to a quarter of a second.

γ Ursæ Majoris.

1812, June 22, Oct. 27, 28, Nov. 3, 5, 6, 21, 24, Dec. 6. 1813, Mar. 17 -	55,440	Mean of 10	55,440
Mar. 21, Apr. 1, 3, 4, 7, 8, 9, 10, 11, 12 - -	54,975	Mean of 20	55,207
Apr. 13, 14, 16, 17, 20, 26, May 22, 26, 28, 29 -	55,489	Mean of 30	55,301
May 31, June 1, 7, 8, 12, 23, 25, 27, July 5, 6 -	55,457	Mean of 40	55,340
July 16, 20, 30, Aug. 7, 20, 31, Sept. 12, 14 - -	54,952	Mean of 48	55,275

This result, like the last, is probably exact to a quarter of a second.

γ Draconis.

1812, June 15, 16, 21, 23, 28, 29, July 6, 7, 8, 9 -	3′,845	Mean of 10	3′,845
July 10, 11, 14, 17, 19, 20, 21, 22, 28, 29 - -	3.779	Mean of 20	3.812
July 30, 31, Aug. 1, 3, 12, 13, 15, 17, 20, 21 - -	3.182	Mean of 30	3.602
Sept. 15, 16, 18, 19, 20, 21, Oct. 1, 3, 5, 8 - -	3.604	Mean of 40	3.603
Oct. 9, 15, 21, 24, 26, 28, 29 31, Nov. 3, 6 - -	3.651	Mean of 50	3.612
Nov. 8, 15, 19, 20, 21, 22, 23, Dec. 5, 6, 8 - -	3.877	Mean of 60	3.656
Dec. 9, 10, 13, 15. 1813, June 22, 24, 25, 26, 27, 28 -	3.471	Mean of 70	3.635
July 5, 6, 9, 10, 11, 12, 13, 16, 17, 18 - -	3.980	Mean of 80	3.674
July 19, 20, 22, 23, 24, 25, 26, 27, 28, 29 - -	3.477	Mean of 90	3.652

This determination of γ Draconis is extremely exact. Its zenith distance has also been determined, with equal accuracy, by 120 observations made with the zenith sector during the years 1811 and 1812: the mean of above 60 observations of 1811 do not differ more than one-tenth of a second from the mean of an equal number in 1812.

	′	″
Mean zenith distance beginning of 1813 by observations of 1811 - - -	2	17.8
By observations in 1812 - - -	2	17.9
Mean zen. distance by zenith sector -	2	17.85
N. P. D. by mural circle - -	38 29	3.65
Sum or Co. latitude - - -	38 31	21.5

η Ursæ Majoris.

1812, June 11, 12, 13, 14, 15, 17, 20, 20, 22, 23, 24 -	57,671	Mean of 10	57,671
June 28, July 6, 7, 8, 9, 11, 15, 18, 20, 28 - -	58,069	Mean of 20	57,870
Aug. 7, 15, 19, 20, 22, Sept. 16, 18, 20, 21, Oct. 4 -	58,138	Mean of 30	57,968
Oct. 5, 7, 8, 24, 28, 30, Nov. 5, 6, 7, 14 - -	58,000	Mean of 40	57,969
Nov. 18, 21, 23, Dec. 5, 8, 23. 1813, May 26, 28, 31, June 1	57,348	Mean of 50	57,845
June 4, 7, 10, 11, 12, 16, 21, 22, 23, 25 - -	58,123	Mean of 60	57,891
June 26, 27, July 5, 6, 9, 16, 18, 19, 20, 22 -	58,057	Mean of 70	57,915
July 24, 27, 28, 29, 30, Aug. 8, 12, 13, 16, 19 -	57,641	Mean of 80	57,881

This result is probably not inferior in precision to the last.

α Persei.

1812, July 7, 9, 17, 20, 21, 28, Aug. 16, 17, Dec. 8, 12	52,970	Mean of 10	52,970
Dec. 13, 14, 28. 1813, Jan. 8, 10, 11, 22, 24, 28, Feb. 5	52,417	Mean of 20	52,693
Feb. 23, March 1, 4, 6, 7, 17, 19, Apr. 11, 12, 13 -	52,468	Mean of 30	52,618
Apr. 14, 18, May 27, 28, June 1, 19, 20, July 6, 15, 17	52,824	Mean of 40	52,670

Exact to a quarter of a second.

Capella.

1812, June 15, 26, 28, July 7, 8, 9, 10, 14, 20, 23 -	20,987	Mean of 10	20,987
July 28, 30, Aug. 13, 16, 17, 19, 21. 1813, Jan. 2, 7, 8	21,009	Mean of 20	20,998
Jan. 10, 11, 16, 22, 24, 27, 28, 31, Feb. 23, Mar. 1 -	20,719	Mean of 30	20,905
Mar. 2, 3, 5, 6, 7, 12, 17, 18, 22, Apr. 8 - -	19,951	Mean of 40	20,667
Apr. 9, 10, 11, 12, 13, 14, 15, 17, 18, 20 - -	21,124	Mean of 50	20,758
Apr. 21, May 28, 29, June 1, 2, 20, July 4, 5, 6, 12 -	19,537	Mean of 60	20,555
July 15, 16, 17, 22, 23, 25, 28, 29, 30, 31 - -	20,258	Mean of 70	20,512
Aug. 3, 5, 6, 8, 9, 10, 11, 13, 15, 17 - -	20,061	Mean of 80	20,468

One of the above series differs a second from the mean of the whole, a circumstance very unusual, but quite unconnected with error of division, I attribute it to want of sufficient care in reading off the microscopes. The mean result may, nevertheless, be depended upon to a quarter of a second.

α Cygni.

1812, Sept. 15, 16, 17, 18, 20, 21, Oct. 2, 3, 4, 5 -	56,846	Mean of 10	56,846
Oct. 8, 10, 13, 14, 16, 22, 23, 24, Nov. 7, 8 - -	56,568	Mean of 20	56,707
Nov. 15, 19, 20, 21, 22, 23, 24, Dec. 8, 9, 10 - -	56,980	Mean of 30	56,798
Dec. 13, 30. 1813, Jan. 8, 9, 10, 16, 22, Feb. 8, 9, 10	57,405	Mean of 40	56,925
Feb. 15, 28, March 6, 11, 12, 16, 20, 26, 27, Apr. 1 -	57,178	Mean of 50	56,975
July 27, 28, 29, 30, Aug. 3, 4, 5, 7, 9, 10 - -	56,665	Mean of 60	56,940
Aug. 11, 12, 13, 15, 17, 19, 20, 21, 22, 23 - -	56,887	Mean of 70	56,947

Result very exact.

α Lyræ.

1812, July 6, 7, 9, 10, 14, 17, 21, 22, 23, 26 -	0,123	Mean of 10	0,123
July 28, 30, 31, Aug. 1, 6, 12, 13, 15, 16, 17 - -	0,514	Mean of 20	0,318
Aug. 18, 19, 20, 21, Sept. 15, 16, 18, 19, Oct. 1, 2 -	0,364	Mean of 30	0,334
Oct. 3, 4, 5, 8, 9, 10, 12, 15, 16, 19 - -	0,299	Mean of 40	0,325
Oct. 21, 24, 26, 28, 29, 31, Nov. 2, 3, 7, 8 -	0,983	Mean of 50	0,457
Nov. 13, 15, 19, 20, 21, 22, 23, Dec. 2, 6, 8 - -	0,857	Mean of 60	0,522
Dec. 9, 10, 12, 13. 1813, Jan. 2, 7, 9, 15, 21, June 22	0,514	Mean of 70	0,521
June 23, 24, 25, 26, 27, 28, July 5, 6, 9, 10 - -	0,501	Mean of 80	0,519
July 11, 12, 13, 16, 17, 18, 19, 23, 24, 25 - -	0,187	Mean of 90	0,482

Castor.

1812, Aug. 13, 19, 21. 1813, Jan. 16, 24, 28, Mar. 5, 7, 10, 16 - -	46″,916	Mean of 10	46″,916
Mar. 18, 23, 31, Apr. 3, 8, 11, 13, 15, 28, June 1 -	46,672	Mean of 20	46,794
Aug. 11, 14, 16, 23, 24, 28, Sept. 2, 6, 7, 10 -	46,596	Mean of 30	46,728

Pollux.

1812, June 13, 15, 19, Aug. 13, 16, 17, 19, Oct. 1, 3, 4	56″,536	Mean of 10	56″,536
Oct. 11. 1813, Jan. 16, 22, 24, Feb. 28, Mar. 5, 10, 13, 18, 23	56,332	Mean of 20	56,434
Mar. 31, Apr. 8, 11, 13, 15, May 28, June 1, 25, July 28, 30	56,313	Mean of 30	56,394
Aug. 3, 5, 11, 16, 22, 23, 24, 25, 28, Sept. 1 -	56,380	Mean of 40	56,341

Exact to a quarter of a second.

β Tauri.

1812, July 20, 22, 30, Aug. 13, 19, 21, Nov. 19, 28. 1813, Jan. 2, 8 - -	43″,650	Mean of 10	43″,650
Jan. 10, 11, 13, 24, 25, 27, 28, 31, Mar. 1, 3 -	43,438	Mean of 20	43,544
Mar. 5, 6, 7, 12, 17, 18, 19, 31, Apr. 9, 10 - -	43,410	Mean of 30	43,499
Apr. 11, 12, 14, July 15, 16, 17, 25, 28, 29, 30 - -	43,903	Mean of 40	43,600
July 31, Aug. 6, 8, 9, 11, 15, 18, 20, 22, 23 -	43,820	Mean of 50	43,663

Exact to less than a quarter of a second.

α Cor. Bor.

1812, June 11, 12, 13, 14, 15, 19, 21, 22, 23, 28 -	55″,441	Mean of 10	55″,441
June 30, July 4, 6, 8, 9, 15, 18, 19, 20, 21 - -	55,295	Mean of 20	55,368
July 22, 26, 28, Aug. 13, 15, 17, 18, 19, Sept. 15, 16	55,316	Mean of 30	55,351
Sept. 19, 20, 21, Oct. 3, 4, 5, 6, 8, 13, 20 - -	56,041	Mean of 40	55,523
Oct. 26, 28, 29, 31, Nov. 20, 21, 22, Dec. 7, 12, 14 -	55,791	Mean of 50	55,577
1813, Jan. 8, 9, 11, May 29, June 7, 8, 10, 12, 16, 18	54,833	Mean of 60	55,453
June 21, 22, 24, 25, 27, 29, July 5, 6, 10, 12 -	55,008	Mean of 70	55,389
July 16, 17, 18, 19, 20, 23, 24, 26, 27, 28 - -	55,487	Mean of 80	55,401

Exact to less than a quarter of a second.

α Arietis.

1812, June 18, 19, 22, 23, 26, 28, July 6, 7, 9, 10 -	37″,380	Mean of 10	37″,380
July 14, 17, 20, 21, 22, Aug. 17, Oct. 24, 25, 26, 28 -	36,263	Mean of 20	36,822
Oct. 29, 31, Nov. 3, 5, 6, 7, 8, 9, 15, 19 - -	35,811	Mean of 30	36,485
Nov. 21, 22, 24, 28, 29, Dec. 6, 7, 8, 13. 1813, Jan. 8	36,532	Mean of 40	36,497
Jan. 9, 16, May 22, June 19, 20, July 4, 5, 6, 11, 16	36,474	Mean of 50	36,492

I mentioned this star as doubtful in my former Catalogue. I believe the discordances were quite accidental, and that the mean result is within a quarter of a second of the truth.

Arcturus.

1812, June 11, 12, 13, 15, 16, 19, 20, 21, 28, 29 -	18″,729	Mean of 10	18″,729
July 4, 6, 7, 8, 9, 11, 15, 18, 20, 26 - -	19,867	Mean of 20	19,298
July 18, 31, Aug. 7, 14, 18, 19, Sept. 20, 21, Oct. 2, 4 -	19,341	Mean of 30	19,312
Oct. 7, 9, 13, 16, 18, 24, 25, 26, Nov. 3, 5 - -	19,225	Mean of 40	19,291
Nov. 6, 7, 14, 18, 21, 22, 23, Dec. 6, 7, 8 - -	19,453	Mean of 50	19,323
Dec. 12, 14, 23. 1813, May 24, 26, 27, 28, 31, June 1, 4	18,761	Mean of 60	19,230
June 7, 8, 10, 11, 12, 16, 21, 22, 25, 26 - -	18,585	Mean of 70	19,137
June 27, July 5, 6, 7, 9, 12, 16, 18, 19, 23 - -	18,653	Mean of 80	19,077

Exact to a quarter of a second, or less.

Aldebaran.

1812, June 23, 28, July 5, 7, 8, 9, 10, 25, 26, 28 -	35″,614	Mean of 10	35″,614
Aug. 14, 16, 17, 19, 21, Sept. 5, Dec. 25, 28. 1813, Jan. 8, 10	35,135	Mean of 20	35,374
Jan. 11, 16, 22, 24, 28, 31, Feb. 23, Mar. 1, 6, 7 -	35,475	Mean of 30	35,407
Mar. 11, 17, 18, 19, 31, Apr. 4, 8, 9, 10, 11 - -	35,390	Mean of 40	35,402
Apr. 12, 14, 15, June 28, July 6, 8, 10, 15, 16, 17 -	35,347	Mean of 50	35,392
July 23, 25, 27, 28, 29, 30	35,142	Mean of 56	35,365

Result very exact.

β Leonis.

1812, June 13, 14, 15, 22, 29, July 6, 8, 9, 11, 12 -	57″,544	Mean of 10	57″,544
July 15, 23, Oct. 2, 3, 4, 6, 7, 14, 15, 22 - -	57,077	Mean of 20	57,310

This star has not been often observed, owing to its proximity in right ascension to γ Urs. Maj.; but the result is probably exact to within 0″,3, as it very seldom happens that the mean of twenty observations differs a quarter of a second from the truth

α Herculis.

1812, July 11, 14, 30, Aug. 3, 7, 15, 17, 18, 20, Sept. 19	13″,835	Mean of 10	1″,3835
Sept. 20, 21, Oct. 3, 4, 8, 9, 29, 31, Nov. 15, 21 -	14,170	Mean of 20	14,002
Nov. 22, 23, Dec. 12. 1813, Jan. 9, 11, 15, July 11, 25, 27, 28 - - -	13,803	Mean of 30	13,936
July 29, 30, Aug. 2, 4, 7, 9, 10, 12, 13, 15 - -	14,223	Mean of 40	14,008
Aug. 17, 18, 19, 23, 24, 30, 31, Sept. 5, 6, 13 -	14,169	Mean of 50	14,040

Result very exact.

α Pegasi.

1812, Oct. 1, 2, 3, 4, 5, 8, 10, 11, 12, 16 - -	51″,880	Mean of 10	51″,880
Oct. 17, 21, 28, Nov. 3, 7, 9, 19, Dec. 5, 7, 12 -	52,060	Mean of 20	51,695

This star is reserved for future examination.

Regulus.

1812, June 12, 13, 15, 19, 20, 22, 23, July 6, 7, 8 -	23,331	Mean of 10	23,331
July 9, 14, 15, 18, 21, 23, 28, Oct. 1, 2, 3 - -	23,205	Mean of 20	23,268
Oct. 4, 6, 12, 14, 15, 19, 20, 23, 27, 28 - -	22,543	Mean of 30	23,026
Nov. 5. 1813, Mar. 7, 10, 12, 13, 17, 18, 22, 23, 26 -	22,236	Mean of 40	22,829
Mar. 29, 31, Apr. 1, 2, 3, 4, 7, 8, 9, 11 - -	22,148	Mean of 50	22,692
Apr. 12, 13, 14, 20, 26, May 24, 28, 31, June 2, 7 -	22,665	Mean of 60	22,688
June 8, 11. July 6, 7, 9 -	22,576	Mean of 65	22,679

Not to be relied on with the same confidence as some others, as one of the above series differs a second from the mean; the result may be erroneous a quarter of a second, but, I think, not much more.

α Ophiuchi.

1812, June 28, 29, July 11, 14, 21, 22, 26, 28, 30, Aug. 3	39,126	Mean of 10	39,126
Aug. 19, 20, Sept. 15, 16, 19, 20, 21, Oct. 3, 8, 9 -	39,949	Mean of 20	39,537
Oct. 28, 31, Nov. 7, 15, 19, 21, 22, 23, Dec. 6, 8 -	40,358	Mean of 30	39,811
Dec. 29. 1813, Jan. 7, 9, 15, June 24, 25, 26, 27, July 5, 6	39,215	Mean of 40	39,662
July 9, 10, 11, 12, 13, 16, 17, 18, 19, 22 - -	38,849	Mean of 50	39,500
July 23, 24, 25, 26, 27, 28, 29, Aug. 2, 4, 5 - -	38,523	Mean of 60	39,339
Aug. 7, 9, 10, 12, 13, 15, 17, 19, 20, 24 - -	38,115	Mean of 70	39,163

Some discordances in the above observations, which I cannot explain, render the result doubtful, and I reserve this star for future examination.

α Aquilæ.

1812, July 26, 30, Aug. 1, 3, 6, 9, 12, 14, 18, 19 -	58,133	Mean of 10	58,133
Sept. 15, 16, 18, 19, 20, 21, Oct. 2, 3, 4, 7 - -	59,381	Mean of 20	58,757
Oct. 8, 10, 13, 15, 16, 19, 21, 23, 24, 27 - -	58,963	Mean of 30	58,826
Oct. 28, 29, Nov. 2, 3, 6, 7, 15, 19, 21, 22 - -	59,705	Mean of 40	59,045
Nov. 23, 24, Dec. 8, 9, 10. 1813, Jan. 21, Feb. 28, Mar. 5, 6, 7 - -	59,626	Mean of 50	59,162
Mar. 12, 20, 26, 27, Apr. 1, 3, July 11, 12, 16, 17 -	58,014	Mean of 60	58,970
July 19, 21, 22, 25, 26, 27, 28, 29, 30, Aug. 1 -	58,280	Mean of 70	58,872
Aug. 2, 3, 4, 7, 10, 11, 12, 13, 15, 16 - -	58,280	Mean of 80	58,799

Procyon.

1812, June 12, 19, July 25, Sept. 15, 16, 17, 18, 19, 20. 1813, Feb. 24 - -	13,999	Mean of 10	13,999
Mar. 3, 6, 8, 17, 22, 29, Apr. 2, 4, 9, 10 - -	14,221	Mean of 20	14,110
Apr. 12, 14, 20, June 2, 7, 8, July 29, Aug. 11, 14, 16	14,795	Mean of 30	14,338
Aug. 17, 23, 24, Sept. 2, 6, 7, 10, 12, 13, 14 -	14,426	Mean of 40	14,360

Probably exact to a quarter of a second.

α Orionis.

1812, July 17, 22, 25, Aug. 5, 13, 14, 16, 17, 19. 1813, Jan. 2 - -	16′,044	Mean of 10	16′,044
Jan. 10, 11, 13, 16, 22, 25, 27, 28, Mar. 2, 11 -	15,486	Mean of 20	15,765
Mar. 18, 22, Apr. 3, 4, 9, 10, 11, 12, 13, 14 -	16,380	Mean of 30	15,970
Apr. 17, 18, 19, 20, May 28, July 8, 16, 17, 28, 29 -	15,849	Mean of 40	15,940
Aug. 6, 8, 11, 13, 18, 19, 20, 23, 24, Sept. 1 -	14,856	Mean of 50	15,793

From the discordance of the last series, I consider this result as doubtful.

α Serpentis.

1812, June 12, 13, 17, 19, 22, 23, 28, 30, July 2, 4 -	38′,187	Mean of 10	38′,187
July 6, 8, 9, 15, 18, 20, 21, 22, 26, 28 - -	39,508	Mean of 20	38,847
Aug. 14, 15, 17, 18, 19, Sept. 15, 16, 18, 21, Oct. 3 -	39,700	Mean of 30	39,132
Oct. 4, 13, 26, Dec. 6, 7, 12. 1813, Jan. 8, 9, 15, June 7	39,282	Mean of 40	39,169
June 8, 10, 13, 16, 18, 21, 22, 24, 25, 27 - -	39,244	Mean of 50	39,184
July 5, 10, 11, 12, 16, 17, 18, 19, 20, 22 - -	39,670	Mean of 60	39,265
July 23, 26, 27, 28, 29, 30, Aug. 2, 7, 12, 15 -	39,240	Mean of 70	39,260

Result very exact.

Polaris.

Mean Day of the Month.	Position of the Telescope on the Instrument.	Number of Observations above the Pole.	Number of Observations below the Pole.	Total Number of Observations.	N. P. D. Jan. 1, 1813.				
1812.					°	′	″		″
June 15	0	3	8	11	1	41	21,79	Mean of 11	21,79
23	30	4	3	7			22,06	Mean of 18	21,89
July 7	10	4	4	8			22,49	Mean of 26	22,08
18	20	6	5	11			22,29	Mean of 37	22,14
Oct. 18	0	10	9	19			21,69	Mean of 56	21,99
Nov. 10	10	10	6	16			21,71	Mean of 72	21,95
Dec. 8	20	10	7	17			21,82	Mean of 89	21,83
1813.									
Apr. 1	30	19	17	36			21,44	Mean of 125	21,72
June 10	0	19	23	42			21,70	Mean of 167	21,72

The mean of more than 200 observations of this star is 1° 41′ 21″,75. The above 167 were selected in preference; the result, however, is the same.

Notwithstanding the great number of observations of this star, there are discordances which render the result doubtful to 0″,25. If the observations with four microscopes, previous to Oct. 1812, be rejected, the mean result with six microscopes will be 1° 41′ 21″,6, which I prefer to the above.

The following Table shews the State of the Standard Catalogue at this present Time, Sept. 1813.*

N. P. D. of Stars for the beginning of the Year 1813.

	Names of Stars.	No. of Obs.	N. P. D. Jan. 1813.			Difference of former Catalogue.	
			°	′	″		
1	Polaris	200	1	41	21,75	+ 0,02	
2	β Urs. min.	90	15	4	48,95	0,00	
3	β Cephei	40	20	5	30,70	+ 0,40	
4	α Urs. maj.	60	27	14	31,46	— 0,04	
5	α Cephei	40	28	12	12,47	+ 0,12	
6	α Cassiop.	40	34	29	22,73	— 0,10	
7	γ Urs. maj.	50	35	15	55,47	+ 0,17	
8	γ Draconis	90	38	29	3,65	— 0,08	
9	η Urs. maj.	80	39	44	57,88	+ 0,08	
10	α Persei	40	40	48	52,67	+ 0,30	
11	Capella	80	44	12	20,47	— 0,49	
12	α Cygni	800	45	22	56,92	= 0,27	
13	α Lyræ	100	51	23	0,43	— 0,29	
14	Castor	30	57	42	46,73	+ 0,16	
15	Pollux	40	61	31	56,34	— 0,23	
16	β Tauri	50	61	34	43,66	+ 0,15	
17	α Androm.	35	61	56	29,61		The same as in former Catalogue,
18	α Cor. Bor.	90	62	38	55,43	— 0,27	and probably true to 0″,5 or less.
19	α Arietis	50	67	25	36,49	— 0,27	
20	Arcturus	80	69	50	19,68	+ 0,04	
21	Aldebaran	56	73	52	35,36	+ 0,18	
22	β Leonis	20	74	22	57,31	+ 0,07	
23	α Herculis	50	75	23	14,04	+ 0,07	
24	α Pegasi	20	75	47	51,70	+ 0,07	
25	Regulus	50	77	7	22,69	— 0,25	
26	α Ophiuchi	70	77	17	39,16	— 0,50	Doubtful to 0″,5.
27	α Aquilæ	100	81	36	58,66	— 0,22	Doubtful.
28	α Orionis	30	82	38	15,72	— 0,24	Doubtful to 0″,5.
29	α Serpentis	70	82	58	39,26	+ 0,39	
301	Procyon	40	84	18	14,36	— 0,36	

‡ The N. P. D. of Procyon in the former Catalogue was 15″,03; this was from a mistake of 1″,0 committed in adding the annual variation, it should have been 14″,03.

* Though the Observations were given to the Society, as by the date of the paper, yet, by the permission of the President and Council, they were extended till the time that they went to the press.

Remarks on the above Observations.

α Lyræ and α Aquilæ having been supposed subject to a sensible parallax, I have, as I mentioned before, reserved them for future examination. The observations which I have already made on these stars, and particularly on α Aquilæ, are not incompatible with this supposition, though I cannot at present venture to decide whether the small discordances I have met with are to be attributed to any regular cause, or are only accidental.

Whenever I speak of the degree of exactness to which any result may be depended upon, I allude only to the mechanical measure given by the instrument. I have every reason to believe, that if two fixed and well defined points could be placed in the plane of the meridian, I could, in a very short time, measure their angular distance to within a tenth of a second; but astronomers must be well aware that the stars are not presented to us in this simple form, and that the sources from which small errors may arise, either in the observations themselves or subsequent computation, are so very numerous, that anomalies will occur even to the most careful observer, which he will in vain endeavour to explain. With respect to the parallax of α Lyræ, I might observe that it is a star so badly defined, and so little adapted for exact observation, that a parallax of half a second would not be easy to determine even with the Greenwich circle.

α Aquilæ is in some respects a better star for observation, but only half its actual parallax would be sensible in declination.

There are several other stars much better adapted for this investigation, even should their distance be supposed more than double, such are Polaris, η Ursæ maj. α Cygni, β Urs. min. and γ Draconis; now in these I have not hitherto found any sensible parallax; occasional discordance has frequently suggested some slight hopes, but these have always been disappointed by continuing the observations. It is, however, useless now to anticipate this subject farther.

Those stars which are in the general Catalogue, but which do not form part of the standard Catalogue, I presume to be exact to the nearest second.

I have not included any star in the standard Catalogue south of the equator, on account of the uncertainty of refraction.

XXXIII. *Observations of the Summer Solstice, 1813, with the Mural Circle, at the Royal Observatory.*

By John Pond, *Esq. Astronomer Royal, F. R. S.*

Read July 8, 1813.

1813.	Barometer.	Therm. In.	Therm. Out.	Refraction.	Observations as given by the Instrument.	Equations for N. P. D.	Equations for Zenith Distance.	Semidiameter of the ☉ by Nautical Almanack.	Reduction to the Solstice.	Solstitial Zenith Distance with Parallax.	Solstitial N. P. D. with Parallax.	Correction for ☉'s Lat.	Solstitial Zenith Distance corrected for ☉'s Lat.	Solstitial N. P. D. corrected for ☉'s Lat.
				′ ″		″	° ′ ″	′ ″	′ ″	° ′ ″	° ′ ″	″	° ′ ″	° ′ ″
June 10	29,57	62	69	30,3	☉ LL 67 14 21,9	— 0,6	— 38 31 22,1	15 46,5	— 26 44,7	28 0 58,9	66 32 20,4	+ 1,0	28 0 59,9	66 32 21,4
11	29,81	64	71	29,6	☉ UL 66 38 18,9	— 0,6	22,1	15 46,5	22 10,2	56,7	18,2	+ 1,0	0 57,7	19,2
12	29,70	64	74	30,1	☉ LL 67 5 48,1	— 0,6	22,1	15 46,4	18 12,1	57,6	19,1	+ 1,0	0 58,6	20,1
13	30,02	61	66	30,9	☉ UL 66 30 36,9	— 0,6	22,1	15 46,3	14 32,4	59,6	21,1	+ 0,9	1 0,5	22,0
15	29,62	60	66	30,1	☉ LL 66 56 2,3	— 0,6	22,1	15 46,1	8 20,7	57,5	19,0	+ 0,7	0 58,2	19,7
21	30,15	57	60	30,0	☉ UL 66 16 9,3	— 0,6	22,1	15 45,8	0 1,9	1 1,1	22,6	— 0,1	1 1,0	22,5
23	30,17	56	59	30,8	☉ LL 66 48 7,4	— 0,6	22,1	15 45,6	0 32,0	0 58,5	20,0	— 0,3	0 58,2	19,7
25	30,18	59	64	29,9	☉ UL 66 18 47,8	— 0,6	22,1	15 45,6	2 41,1	1 0,1	21,6	— 0,3	0 59,8	21,3
27	30,07	64	70	30,2	☉ LL 66 54 6,5	— 0,6	22,1	15 45,6	6 29,3	0 59,7	21,2	— 0,2	0 59,5	21,0
28	29,94	61	67	29,7	☉ UL 66 25 9,0	— 0,6	22,1	15 45,6	9 0,4	1 1,8	23,3	— 0,1	1 1,7	23,2
29	29,75	64	74	30,0	☉ LL 66 59 32,2	— 0,6	22,1	15 45,5	11 56,1	0 58,5	20,0	0,0	0 58,5	20,0
July 1	29,67	62	65	29,6	☉ UL 66 35 7,1	— 0,6	22,1	15 45,5	19 0,7	0 59,4	20,9	+ 0,3	0 59,7	21,2
2	29,75	60	62	31,8	☉ LL 67 10 44,8	— 0,6	22,1	15 45,5	23 9,5	59,5	21,0	+ 0,4	0 59,9	21,5
					Mean of 13 Observations -					28 0 59,1	66 32 20,6		28 0 59,5	66 32 21,0
					Parallax — 4″,0 Nutation — 6″,5 =					— 10,5	— 10,5		— 10,5	— 10,5
										28 0 48,6	66 32 10,1		28 0 49,0	66 32 10,5
					Mean Obliquity - -									23 27 49,5
					* Mean Obliquity at Summer Solstice, 1812									23 27 50,5
					Mean of Two Observations or Mean Obliquity, Jan. 1, 1813 -									23 27 50,0

* I avail myself of this opportunity of correcting a very small error that was made in computing the summer solstice of 1812. The correction for the sun's latitude should have been 0″,6 instead of 0″,9, and should have been applied with the contrary sign. The obliquity thus corrected will be 23° 27′ 50″,5.

PRESENTS

RECEIVED BY THE

ROYAL SOCIETY,

From November 1812 *to July* 1813,

WITH THE

NAMES OF THE DONORS.

1812.	PRESENTS.	DONORS.
Nov. 5.	A Description of the Collection of ancient Marbles in the British Museum with Engravings. Part I. London, 1812. 4°	The Trustees of the British Museum.
	A Treatise on Theatres, by George Saunders. London, 1790. 4°	Mr. George Saunders.
	Tracts on Mathematical and Philosophical Subjects, by Charles Hutton, LL. D. and F. R. S. London, 1812. 3 Vols. 8°	Dr. Charles Hutton.
	The Elements of Botany, or Outlines of the Natural History of Vegetables, by Benjamin Smith Barton, M. D. Philadelphia, 1812. 8°	Dr. Benj. Smith Barton.
	Elements of Chemical Philosophy, by Sir Humphry Davy. Part I. Vol. I. London, 1812. 8°	Sir Humphry Davy.
	Essai sur la Rage, par M. J. F. A. Lalouette. 8°	Mr. J. F. A. Lalouette.
	Institutions de Physique, par B. G. Sage. Paris, 1811. 3 Vols. 8°	Mr. B. G. Sage.
	Of the Circle and the infinite Incommensurability of its Area to the Square of its Diameter, with a Demonstration of Dr. Matthew Stewart's 42d Proposition, by James Glennie, Esq. F. R. S. 4°	James Glennie, Esq.
	The Speech of His Royal Highness the Duke of Sussex in the House of Lords, April 21, 1812. 4°	The Editor.
	An Account of a Case of Recovery, after an extraordinary Accident, in which the Shaft of a Chaise had been forced through the Thorax, by William Maiden. London, 1812. 4°	Mr. William Maiden.
	Eloge Historique de M. Sabatier, par M. Percy. Paris, 1812. 8°	Mr. Percy.
	Della Cometa del 1811, osservata nella Specola di Palermo, par Gius. Piazzi. 4°	Mr. Piazzi.

1813.	PRESENTS.	DONORS.
Nov. 5.	Observations on Dr. Sturges Pamphlet respecting Non-residence of the Clergy, in a Letter from T. B. Howell, Esq. to Mr. Baron Maseres. London, 1803. 8°	T. B. Howell, Esq.
	The Philosophical Magazine from June to October, 1812. 8°	Mr. A. Tilloch.
	The Critical Review from June to October, 1812. 8°	The Proprietor.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 148 to 152. 8°	Mr. W. Nicholson.
	The European Magazine from January to October, 1812.	The Editor.
12.	A Catalogue of the Archiepiscopal Manuscripts in the Library at Lambeth Palace, with an Account of the Archiepiscopal Registers and other Records there preserved. fol.	The Archbishop of Canterbury.
	Illustrations of the Lotus of Antiquity, by R. Duppa, LL.B. London, 1813. 4°	R. Duppa, Esq.
	Kongl Vetenskaps Academiens nya Handlingar, Tom. XXX, XXXI, XXXII, for the Years 1809, 1810, and 1811. 8°	The Royal Academy of Sciences of Stockholm.
	A Sermon preached before the incorporated Society for the Propagation of the Gospel in Foreign Parts, Feb. 1, 1812, by Samuel Lord Bishop of Carlisle. London, 1812. 4°	The Bishop of Carlisle.
	De l'Opération de la Cataracte, par A. E. Tartra. Paris, 1812. 4°	Mr. A. E. Tartra.
	Quelques Idées nouvelles sur le Système de l'Univers, par Guil. Ant. Marechal. Paris, 1810. 8°	Mr. G. A. Marechal.
	Mon Opinion sur la Formation des Aerolithes, par Guil. Ant. Marechal. Paris, 1812. 8°	———
19.	Abstract of Answers and Returns made pursuant to an Act (51 George III.) for taking an Account of the Population of Great Britain, and of the Increase and Diminution thereof, 1811. fol.	The Speaker of the House of Commons.
	Comparaison des Organes de la Mastication des Orthopterès, par M. Marcel de Serres. 4°	Mr. Marcel de Serres.
	De l'Odorat et des Organes qui paroissent en être le Siége chez les Orthoptères, par M. Marcel de Serres. 4°	———
	Supplement aux Institutions de Physique, par B. G. Sage. Paris, 1812. 8°	Mr. B. G. Sage.
	Flora Lapponica, Auctore Geor. Wahlenberg. Berlin, 1812. 8°	Mr. G. Wahlenberg.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 153.	Mr. W. Nicholson.
Dec. 10.	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 154.	———
	The Philosophical Magazine for November.	Mr. A. Tilloch.
	The Critical Review for November.	The Proprietor.
	The European Magazine for November.	The Editor.

1812.	PRESENTS.	DONORS.
Dec. 17.	Sui Mezzi immediate di estirpare la Povertà in Sicilia, per D. Giuseppe Emmanuale Ortolani. Palermo, 1812. 4°	Mr. Jos. Emmanuel Ortolani.
24.	Reports from the Commissioners of Public Records, with an Account of their Proceedings from 1800 to 1812. fol.	The Commissioners of Public Records.
	Rotuli Hundredorum Temp Hen. III. et Edw. I. In Turr. Lond. et in Curia Receptæ Scaccarii Westm. asservati. Vol. I. fol.	———
	Inquisitionum ad Capellam Domini Regis retornatarum quæ in publicis Archivis Scotiæ adhuc servantur Abbreviatio. Vol. I. II. fol.	———
	A Catalogue of the Harleian Manuscripts in the British Museum, with Indexes of Persons, Places, and Matters. Vol. IV. fol.	———
1813.		
Jan. 14.	The Critical Review, for December, 1812.	The Proprietor.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 154.	Mr. W. Nicholson.
	The Philosophical Magazine for December, 1812.	Mr. A. Tilloch.
	Annals of Philosophy, No. I.	Dr. Thomas Thomson.
	The European Magazine for December, 1812.	The Editor.
28.	Fauna Orcadensis, or the Natural History of the Quadrupeds, Birds, Reptiles, and Fishes of Orkney and Shetland, by the Rev. Geo. Low, &c. Edinburgh, 1813. 4°	Dr. Wm. Elford Leach.
	Medico-Chirurgical Transactions, published by the Medical and Chirurgical Society of London. Vol. III. London, 1812. 8°	The Medical and Chirurgical Society of London.
	Transactions of the Society instituted at London for the Encouragement of Arts, Manufactures, and Commerce. Vol. 30. 8°	Society for the Encouragement of Arts, Manufactures and Commerce.
Feb. 4.	Physiological Reflections on the destructive Operation of spirituous and fermented Liquors on the Animal System, by Thomas Forster. London, 1812. 8°	Thomas Forster, Esq.
	Annals of Philosophy, No. II.	Dr. Thomas Thomson.
	The Critical Review for January.	The Proprietor.
	The European Magazine for January.	The Editor.
	The Philosophical Magazine for January.	Mr. A. Tilloch.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 155 and 157.	Mr. W. Nicholson.
	Biblia Sacra ita exacte translata ut statim videatur quid refert unaquæque vox textûs.	The Abbé Devay.
11.	A Voyage to the South Seas, undertaken by Command of His Majesty for the Purpose of conveying the Bread Fruit Tree to the West Indies in His Majesty's Ship Bounty, commanded by Lieut. Wm. Bligh. London, 1792. 4°	Rear Admiral Bligh.
	A Chart for the Purpose of exhibiting the Tract of His Majesty's Ship Director, in a Voyage to St. Helena and return to England, by Capt. Wm. Bligh, 1800. 1 Sheet.	———

1813.	PRESENTS.	DONORS.
Feb. 25.	A Survey of the Bay of Dublin, by Capt. Wm. Bligh, 1803. 2 Sheets.	Rear Admiral Bligh.
	Chart of the Peninsula and Islands beyond the River Ganges, by James Horsburgh, Esq. 1813. 1 Sheet.	James Horsburgh, Esq.
Mar. 4.	An Introduction to Medical Literature, including a System of practical Nosology, by Thomas Young, M. D. &c. London, 1813. 8°	Dr. Young.
	The Critical Review for February.	The Proprietor.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 158.	Mr. W. Nicholson.
	Annals of Philosophy, No. 3.	Dr. Thomas Thomson.
	The Philosophical Magazine for February.	Mr. A. Tilloch.
	The European Magazine for February.	The Editor.
Mar. 11.	Appendix to the Doctrine of Life Annuities and Assurances, containing a Paper read before the Royal Society on a new Method of calculating the Value of Life Annuities, by Francis Baily. London, 1813. 8°	Francis Baily, Esq.
Apr. 1.	An Examination of the Imposture of Ann Moore, called the Fasting Woman of Tutbury, illustrated by Remarks on other Cases of real and pretended Abstinence, by Alexander Henderson, M. D. London, 1813. 8°	Dr. Henderson.
	Bibliothecæ quam vir doctus et admodum reverendus Daniel Williams, S. T. P. bono publico legavit Catalogus. Londini, 1801. 8°	The Trustees of Dr. D. Williams's Library.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 159.	Mr. W. Nicholson.
	Annals of Philosophy, No. 4.	Dr. Thomas Thomson.
	The Philosophical Magazine for March.	Mr. A. Tilloch.
	The Critical Review for March.	The Proprietor.
8.	Transactions of the Royal Society of Edinburgh. Vol. VI. Part II. Edinburgh, 1809. 4°	The Royal Society of Edinburgh.
29.	An Account of what appeared on opening the Coffin of King Charles I. in the Vault of King Henry VIII. in St. George's Chapel at Windsor, on the 1st of April, 1813, by Sir Henry Halford, Bart. F. R. S. London, 1813. 4°	Sir Henry Halford, Bart.
	Bija Ganita, or the Algebra of the Hindus, by Edward Strachey, Esq.	Edward Strachey, Esq.
	Anatomy of the Heart, Cranium, and Brain, adapted to the Purposes of the medical and surgical Practitioner, and Observations on the Laws of Life and Sensation, by Alex. Ramsay, M. D. 2d Edition. Edinburgh, 1813. 4°	Dr. Alex. Ramsay.
	A Series of Plates of the Heart, Cranium, and Brain, in Imitation of Dissections, by Alexander Ramsay, M. D. 2d Edition. Edinburgh, 1813. 4°	
	A Treatise on new Philosophical Instruments for various Purposes in the Arts and Sciences, with Experiments on Light and Colours, by David Brewster, LL. D. &c. Edinburgh, 1813. 8°	Dr. David Brewster.

1813.	PRESENTS.	DONORS.
May 6.	The European Magazine for March.	The Editor.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 161.	Mr. W. Nicholson.
	Annals of Philosophy, No. 5.	Dr. Thomas Thomson.
	The Philosophical Magazine for April.	Mr. A. Tilloch.
	The Critical Review for April.	The Proprietor.
	The European Magazine for April.	The Editor.
20.	The Transactions of the Linnean Society of London. Vol. XI. Part I. London, 1813. 4°	The Linnean Society.
	A short Account of Experiments and Instruments depending on the Relations of Air to Heat and Moisture, by John Leslie, F. R. S. Ed. Edinburgh, 1813. 8°	John Leslie, Esq.
	Reports of the late John Smeaton, F. R. S. made on various Occasions in the Course of his Employment as a Civil Engineer. London, 1812. 3 Vols. 4°	The Society of Civil Engineers.
27.	Travels in Sweden during the Autumn of 1812, by Thomas Thomson, M.D. London, 1813. 4°	Dr. Thomas Thomson.
	A Treatise on plane and spherical Trigonometry, by Robert Woodhouse, A. M. Cambridge, 1813. 8°	Robert Woodhouse, Esq.
June 3.	A Topographical Dictionary of Scotland, and the Islands in the British Seas, &c. by Nicholas Carlisle, Sec. A. S. London, 1813. 2 Vols. 4°	Nicholas Carlisle, Esq.
	Notice sur la Vie et les Travaux de M. Maskelyne, lue à la Séance publique de l'Institut National de France de 4 Janvier, 1813, par M. le Chevalier Delambre, Secrétaire perpétuel. Londres, 1813. 4°	Mrs. Maskelyne.
	Tableau méthodique des Espèces minérales, seconde partie, par J. A. H. Lucas. Paris, 1806. 8°	Mr. J. A. H. Lucas.
	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 160 and 162.	Mr. W. Nicholson.
	Annals of Philosophy. No. 6.	Dr. Thomas Thomson.
	The Philosophical Magazine for May.	Mr. A. Tilloch.
	The Critical Review for May.	The Proprietor.
	The European Magazine for May.	The Editor.
24.	A Catalogue of the Library of the London Institution. London, 1813. 8°	The Managers of the London Institution.
July 1.	A Journal of Natural Philosophy, Chemistry, and the Arts. No. 163.	Mr. W. Nicholson.
	The Philosophical Magazine for June.	Mr. A. Tilloch.
	The Critical Review for June.	The Proprietor.
8.	Théorie analytique des Probabilités, par M. le Comte Laplace. Paris, 1812. 4°	The Count Laplace.
	Exposition du Système du Monde, par M. le Comte Laplace. 4th Edition. Paris, 1813. 4°	
	Expériences sur la Principe de la Vie, notamment sur celui des Mouvemens du Cœur, et sur le Siége de ce Principe, par M. Le Gallois. Paris, 1812. 8°	Mr. Le Gallois.

1813.	PRESENTS.	DONORS.
July 8.	Opuscules de Physique, par B. G. Sage. Paris, 1813. 8°	Mr. B. G. Sage.
	Recherches sur la Chaleur developpée dans la Combustion, et dans la Condensation des Vapeurs, par le Comte de Rumford, F. R. S. Paris, 1813. 8°	Count Rumford.
	Recherches sur les Bois et le Charbon, par le Comte de Rumford. Paris, 1813. 8°	———
	Annals of Philosophy. No. 7.	Dr. Thomas Thomson.
	The European Magazine for June.	The Editor.
	Mémoire explicatif sur la Sphère Caucasienne, et spécialement sur le Zodiaque, par C. G. S. Paris, 1813. 4°	The Author.
	Le Zodiaque expliqué, ou Recherches sur l'Origine et la Signification des Constellations de la Sphère Grecque, traduit du Suedois. Paris, 1809. 8°	The Author.
	Tables Astronomiques publiées par le Bureau des Longitudes de France—et Tables de la Lune par M. Burckhardt. Paris, 1813. 4°	M. Burckhardt.

INDEX

TO THE

PHILOSOPHICAL TRANSACTIONS

FOR THE YEAR 1813.

A

page

Acid, liquid fluoric, discovered by Scheele, description of it, - - - - - - 265

———— an account of some experiments undertaken to electrize it, - - - - - 270

Acids, remarks on their influence upon the composition of the urine, - - - - - - - 213

Alcohol, remarks on the state in which it exists in fermented liquors, - - - - - - 82

Alcohol of sulphur, or *Sulphuret of carbon*, experiments on it, 171

———— preparation and general properties of it, 174

———— experiments to ascertain whether hydrogen be present in it, - - - - - 177

———— experiments to ascertain the presence of carbon in it, - - - - - 181

———— particulars respecting the mode in which the proportions of sulphur and carbon in it are established, 188

Ancon, a term applied to a new variety in the breeds of sheep, 93

Ardea Argala, a description of its solvent glands and gizzard, 77

Artery, branchial, observations on its structure - 234

Austin, Rev. Gilbert. On a new construction of a condenser and air pump, - - - 138

Azote and chlorine, a new substance formed by their combination, - - - - - - 2

B

Babington, Dr. His letter to Sir Everard Home on the subject of concretions of olive oil and mucus found in the human intestines, - - - - - 150

INDEX.

page

Bakerian Lecture, on the elementary particles of certain crystals, - - - - - - - 51

BERZELIUS, J. M. D. His experiments on the alcohol of sulphur, or sulphuret of carbon, - - - 171

———————— His appendix to a paper on the alcohol of sulphur, or sulphuret of carbon, - - 188

Bile, has a power of converting animal substance into fat, 153

BLAGDEN, SIR CHARLES. His appendix to Mr. WARE's paper on vision, - - - - - 110

BRANDE, WILLIAM THOMAS, Esq. His additional remarks on the state in which alcohol exists in fermented liquors, 82

———————— His additional observations on the effects of magnesia in preventing an increased formation of uric acid, with remarks on the influence of acids upon the composition of the urine, - - - 213

———————— His experiments to ascertain whether an admixture of bile had any effect upon the process of converting animal substance into fat, - 154

Brentford, Middlesex, an account of some organic remains found near that place, - - - - 181

BREWSTER, DAVID, LL.D. On some properties of light - 101

C

Cassowary, long-legged, from New South Wales, description of its solvent glands and gizzard, - - 79

Cassegrainian telescope, its light compared with that of the Gregorian, - - - - - 206

Casuarius Emu, description of its solvent glands and gizzard, 78

Chemical combination, observations on the atomic doctrine of it, 190

Chlorine and azote, a new substance formed by their combination, - - - - - - 2

Chromate of lead, on the double refraction of it, - 105

CLANNY, WILLIAM REID, M. D. On the means of procuring a steady light in coal mines without the danger of explosion, - - - - - 200

Concave glasses, the bad effects resulting from the use of them described, - - - - - 88

Condenser and air pump, on a new construction of them - 138

COTES's *theorem*, a remarkable application of it, - - 8

Cryophorus, description of an instrument so called, - 78

Crystalline humour, indispensibly necessary to enable the eye to see at different distances, - - - 44

INDEX.

page

Crystals, on the elementary particles of certain, - 51
——— on the existence of two dispersive powers in all doubly refracting, - - - - 107
Cube, origin of its form in crystallography, - - 59

D

DAVY, SIR HUMPHRY, LL.D. On a new detonating compound, - - - - - 1
——— Some further observations on a new detonating substance, - - - 242
——— His experiments and observations on the substances produced in different chemical processes on fluor spar, - - - 269
Diamond, on substances which possess a higher refractive power, - - - - 106

E

Elephant, organic remains of that animal found near Brentford, Middlesex, - - - - 134

F

Fat, on its formation in the intestines of living animals, - 146
Fluor spar, some experiments and observations on the substances produced in different chemical processes on it, - 263
Fluoric combinations, the different hypotheses which may be formed on their nature, - - - 267
——— *principle*, various attempts to procure it in a pure form - - - - - 275
Fluoric compounds, a peculiar substance exists in them, which it is difficult to examine in a pure form, - 278
Fluorine, a name given to a peculiar substance existing in fluoric compounds, - - - 278
Freezing at a distance, on a method of, - - 71

G

Gas, silicated fluoric acid, discovered by SCHEELE, description of it, - - - - - 265
——— *fluo-boric acid*, discovered by GAY LUSSAC and THENARD, description of it, - - - 266
Glands, bronchial, on the black colouring matter of them, - 159
——— *gastric*, experiments to ascertain the coagulating power of their secretion, - - - - 96

page

H

HERSCHEL, J. F. W. Esq. On a remarkable application of Cotes's theorem, - - - - - 8
Hippopotamus, organic remains of that animal found near Brentford, Middlesex, - - - 135
HOME, SIR EVERARD, BART. His description of the solvent glands and gizzards of the Ardea Argala, the Casuarius Emu, and the long-legged Casowary from New South Wales, - - - - - 77
——— His experiments to ascertain the coagulating power of the secretion of the gastric glands, 96
——— On the tusks of the narwhale, 126
——— On the formation of fat in the intestines of living animals, - - - 146
——— His additions to an account of the anatomy of the Squalus Maximus contained in a former paper; with observations on the structure of the branchial artery, - - - - - 227
HOOKE, DR. His account of the formation of certain crystals referred to, - - - - 55
HUMPHREYS, COL. DAVID. On a new variety in the breeds of sheep, - - - - - 88

K

KATER, CAPTAIN HENRY. On the light of the Cassegrainian telescope, compared with that of the Gregorian, - 206

L

Lamp, description of one to be used in coal mines, - 205
Lecture, the Bakerian, on the elementary particles of certain crystals, - - - - - 51
Light, on some properties of it, - - - 101
Lungs, on the colouring matter of the black spots on them, 159

M

Magnesia, additional observations on its effects in preventing an increased formation of uric acid, - - 213
MARCET, ALEXANDER, M. D. His experiments on the alcohol of sulphur, or sulphuret of carbon, - - 171
——— His experiments on the production of cold by the evaporation of the sulphuret of carbon, - - - - - 252

page

Micrometer, description of one with a single lens, - 119
Mines, coal, on the means of procuring a steady light in them without the danger of explosion, - - - 200
——— explosions in, a few of the most considerable in the neighbourhood of Durham recorded, - - 201

N

Narwhale, on the tusks of that animal, - - 126
Near sightedness usually commences between the ages of ten and eighteen, - - - - 81
——— affects the higher classes of society more than the lower, - - - - - - *ib.*
——— sometimes occasioned by an evident change in the spherical figure of the cornea, - - 84
——— seldom alike in the two eyes, - - 85
——— has no dependence on the greater or smaller degree of convexity possessed by the cornea, - - 41
——— not necessarily occasioned by a morbid protrusion of the whole eye, - - - *ib.*

O

Observations of some principal fixed stars, results of them, 285
——— β Ursæ Minoris, - - - *ib.*
——— β Cephei, - - - - 286
——— α Ursæ Major. - - - - *ib.*
——— α Cephei, - - - - *ib.*
——— α Cassiop. - - - - - 287
——— γ Ursæ Major. - - - *ib.*
——— γ Draconis, - - - - 288
——— η Ursæ Major. - - - 289
——— α Persei, - - - - *ib.*
——— Capella, - - - - 290
——— α Cygni, - - - - - 291
——— α Lyræ, - - - - *ib.*
——— Castor, - - - - 292
——— Pollux, - - - - *ib.*
——— β Tauri, - - - - *ib.*
——— α Cor. Bor. - - - - 293
——— α Arietis, - - - - *ib.*
——— Arcturus, - - - - 294
——— Aldebaran, - - - *ib.*
——— β Leonis, - - - 295

page
——— α Herculis, - - - 295
——— α Pegasi, - - - - ib.
——— Regulus, - - - 296
——— α Ophiuchi, - - - 297
——— α Aquilæ, - - - 298
——— Procyon, - - - ib.
——— α Orionis, - - - 299
——— α Serpentis, - - - ib.
——— Polaris, - - - 300
Octohedron, the most common form of crystallized bodies, - 51
Olive oil, concretions of it found in the human intestines, 150

P

PEARSON, GEORGE, M.D. On the colouring matter of the black bronchial glands, and of the black spots of the lungs, 159
POND, JOHN, Esq. *Astronomer Royal*. His observation of the summer solstice of 1812, at the Royal Observatory, - 27
——— His catalogue of north polar distances of some of the principal fixed stars, - 75
——— His observation of the winter solstice of 1812, with the mural circle at Greenwich, 123
——— His catalogue of north polar distances of eighty-four of the principal fixed stars, deduced from observations made with the mural circle at the Royal Observatory, - - - 280
——— Observations of the summer solstice of 1813, with the mural circle, at the Royal Observatory, - - - - 304
PRECHTL, M. His theory respecting various crystalline forms referred to, - - - - 62
Prisms, hexagonal, origin of their form in crystallography, 59

R

Rhea Americana, description of its solvent glands and gizzard, 80
Rhomboid, obtuse, origin of its form in crystallography, - 57

S

Sheep, on a new variety in their breed, - - 88
Sight, observations relative to the near and distant sight of different persons, - - - - 91
SMITHSON, JAMES, Esq. On a substance from the elm tree called ulmin, - - - - 64

INDEX.

page

SMITHSON, JAMES, ESQ. On a saline substance from Mount Vesuvius, - - - - 256

Solstice, summer, observation of that of 1812, at the Royal Observatory, - - - - - 27

——— observation of that of 1813, at the Royal Observatory, - - - - - 298

——— *winter*, observation of that of 1812, with the mural circle at Greenwich, - - - - 123

Spheres, perfect, supposed to be the elementary particles of fluor spar and such other substances as assume the octohedral and tetrahedral forms, - - - - 54

Squalus Maximus, additions to an account of its anatomy contained in a former paper, - - - - 227

Stars, principal fixed, catalogue of north polar distances of some of them. - - - - - 75

——— ———, catalogue of north polar distances of eighty four of them, - - - - 281

Struthio Camelus, description of its solvent glands and gizzard, 80

Sulphuret of carbon, or *Alcohol of sulphur*, experiments on it, 171

——— ——— experiments to determine the proportions of sulphur and carbon contained in it, - - - 182

——— ——— comparison of the proportions of sulphur, and carbon in it, with those which might be inferred from the laws of determinate proportions, - - - 189

——— ——— experiments on its combination with the the alkalies, the earths, and the metallic oxyds, - 192

——— ——— experiments to determine the nature of a particular substance produced by the action of nitro-muriatic acid upon it, - - - - 195

——— ——— experiments on the production of cold by the evaporation of it, - - - - 252

T

Tetrahedron and octohedron, relation of these solids to each other in crystallography, - - - - 51

TRIMMER, MR. WILLIAM KIRBY. His account of some organic remains found near Brentford, Middlesex, - 131

TYCHO L. TYCHONIUS. His account of a narwhale's skull, published in 1706, referred to, - - - 127

U

Ulmin, on a substance from the elm tree, so called, - 64

INDEX.

page

Ulmin, *English*, experiments on it, - - 68
Vesuvius, *Mount*, on a saline substance from it, - 256
Vision, an appendix to Mr. WARE's paper on that subject, 110
Vitreous humour, its absorption probably the cause of amended vision in old persons, - - - 42

W

WARE, JAMES, ESQ. His observations relative to the near and distant sight of different persons, - - 31
Wires, on a method of drawing them extremely fine, - 114
WOLLASTON, WILLIAM HYDE, M.D. Bakerian lecture, on the elementary particles of certain crystals, - - 51
——— On a method of freezing at a distance, - - - 71
——— Method of drawing extremely fine wires, - - 114
——— His description of a single lens micrometer, - - - 119

CORRIGENDA.

Page 175, line 13, *for* one-fourth—*read* one-third
177, —— 17, *for* preponderates over that of either the olive oil or phosphorus—*read* be exceeded by that of the alcohol of sulphur.
246, —— 10, *after* attraction of—*add* the azote of
——— 23, *for* produced—*read* produced,—and dele the comma after portion
249, —— 11, *for* 261.—*read* 268.
264, —— 18, *for* the other—*read* the other of which
268, —— 15, *for* bases,—*read* basis
273, —— 11, *for* metal, positively electrified—*read* metal positively electrified.

From the Press of
W. BULMER & Co.
Cleveland-row, St. James's,
London.